

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



38

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/44, 31/47, 31/50, 31/505, 31/495, 31/53, 31/535, C07D 213/02, 215/00, 217/00, 253/06, 237/06, 239/34, 239/72		A1	(11) International Publication Number: WO 99/01130 (43) International Publication Date: 14 January 1999 (14.01.99)
(21) International Application Number: PCT/US98/13809 (22) International Filing Date: 1 July 1998 (01.07.98) (30) Priority Data: 60/051,592 2 July 1997 (02.07.97) US (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM CORPORATION [US/US]; One Franklin Plaza, Philadelphia, PA 19103 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): ADAMS, Jerry, L. [US/US]; 611 Forest Road, Wayne, PA 19103 (US). HALL, Ralph, F. [US/US]; 1311 Prospect Hill Road, Villanova, PA 19085 (US). (74) Agents: DINNER, Dara, L. et al.; SmithKline Beecham Corporation, Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).			(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CZ, EE, GE, HU, ID, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: NOVEL SUBSTITUTED IMIDAZOLE COMPOUNDS (57) Abstract Novel 1,4,5-substituted imidazole compounds and compositions for use in therapy as CSBP/p38 kinase inhibitors.			
<p>Atty. Docket No. 3015/6/US Serial No. 10/021,780 Anantanarayan et al. Reference 38 of 77</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
DJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

NOVEL SUBSTITUTED IMIDAZOLE COMPOUNDS

5

FIELD OF THE INVENTION

This invention relates to a novel group of imidazole compounds, processes for the preparation thereof, the use thereof in treating CSBP/p38 mediated diseases and pharmaceutical compositions for use in such therapy.

10

BACKGROUND OF THE INVENTION

Intracellular signal transduction is the means by which cells respond to extracellular stimuli. Regardless of the nature of the cell surface receptor (e. g. protein tyrosine kinase or seven-transmembrane G-protein coupled), protein kinases and phosphatases along with phospholipases are the essential machinery by which the signal is further transmitted within the cell [Marshall, J. C. Cell , 80, 179-278 (1995)]. Protein kinases can be categorized into five classes with the two major classes being, tyrosine kinases and serine / threonine kinases depending upon whether the enzyme phosphorylates its substrate(s) on specific tyrosine(s) or serine / threonine(s) residues [Hunter, T., Methods in Enzymology (Protein Kinase Classification) p: 3, Hunter, T.; Sefton, B. M.; eds. vol. 200, Academic Press; San Diego, 1991].

For most biological responses, multiple intracellular kinases are involved and an individual kinase can be involved in more than one signaling event. These kinases are often cytosolic and can translocate to the nucleus or the ribosomes where they can affect transcriptional and translational events, respectively. The involvement of kinases in transcriptional control is presently much better understood than their effect on translation as illustrated by the studies on growth factor induced signal transduction involving MAP/ERK kinase [Marshall, C. J. Cell, 80, 179 (1995); Herskowitz, I. Cell, 80, 187 (1995); Hunter, T. Cell, 80, 225 (1995); Seger, R., and Krebs, E. G. FASEB J., 726-735 (1995)].

While many signaling pathways are part of cell homeostasis, numerous cytokines (e.g., IL-1 and TNF) and certain other mediators of inflammation (e.g., COX-2, and iNOS) are produced only as a response to stress signals such as bacterial lipopolysaccharide (LPS). The first indications suggesting that the signal transduction pathway leading to LPS-induced cytokine biosynthesis involved protein kinases came from studies of Weinstein [Weinstein, *et al.*, J. Immunol. 151,

3829(1993)] but the specific protein kinases involved were not identified. Working from a similar perspective, Han [Han, *et al.*, Science **265**, 808(1994)] identified murine p38 as a kinase which is tyrosine phosphorylated in response to LPS. Definitive proof of the involvement of the p38 kinase in LPS-stimulated signal transduction pathway leading to the initiation of proinflammatory cytokine biosynthesis was provided by the independent discovery of p38 kinase by Lee [Lee; *et al.*, Nature, **372**, 739(1994)] as the molecular target for a novel class of anti-inflammatory agents. The discovery of p38 (termed by Lee as CSBP 1 and 2) provided a mechanism of action of a class of anti-inflammatory compounds for which SK&F 86002 was the prototypic example. These compounds inhibited IL-1 and TNF synthesis in human monocytes at concentrations in the low mM range [Lee, *et al.*, Int. J. Immunopharmac. **10**(7), 835(1988)] and exhibited activity in animal models which are refractory to cyclooxygenase inhibitors [Lee; *et al.*, Annals N. Y. Acad. Sci., **696**, 149(1993)].

MITOGEN AND STRESS ACTIVATED PROTEIN KINASE CASCADES




Figure 1

It is now firmly established that CSBP/p38 is a one of several kinases involved in a stress-response signal transduction pathway which is parallel to and largely independent of the analogous mitogen-activated protein kinase (MAP) kinase cascade

(Figure 1). Stress signals, including LPS, pro-inflammatory cytokines, oxidants, UV light and osmotic stress, activate kinases upstream from CSBP/p38 which in turn phosphorylate CSBP/p38 at threonine 180 and tyrosine 182 resulting in CSBP/p38 activation. MAPKAP kinase-2 and MAPKAP kinase-3 have been identified as downstream substrates of CSBP/p38 which in turn phosphorylate heat shock protein Hsp 27 (Figure 2). It is not yet known whether MAPKAP-2, MAPKAP-3, Mnk1 or Mnk2 are involved in cytokine biosynthesis or alternatively that inhibitors of CSBP/p38 kinase might regulate cytokine biosynthesis by blocking a yet unidentified substrate downstream from CSBP/p38 [Cohen, P. *Trends Cell Biol.*, 353-361(1997)].

p38 Kinase Pathway




Figure 2

What is known, however, is that in addition to inhibiting IL-1 and TNF, CSBP/p38 kinase inhibitors (SK&F 86002 and SB 203580) also decrease the synthesis of a wide variety of pro-inflammatory proteins including, IL-6, IL-8, GM-CSF and COX-2. Inhibitors of CSBP/p38 kinase have also been shown to suppress the TNF-induced expression of VCAM-1 on endothelial cells, the TNF-induced phosphorylation and activation of cytosolic PLA2 and the IL-1-stimulated synthesis of collagenase and

stromelysin. These and additional data demonstrate that CSBP/p38 is involved not only cytokine synthesis, but also in cytokine signaling [CSBP/P38 kinase reviewed in Cohen, P. Trends Cell Biol., 353-361(1997)].

Interleukin-1 (IL-1) and Tumor Necrosis Factor (TNF) are biological substances produced by a variety of cells, such as monocytes or macrophages. IL-1 has been demonstrated to mediate a variety of biological activities thought to be important in immunoregulation and other physiological conditions such as inflammation [See, e.g., Dinarello et al., *Rev. Infect. Disease*, 6, 51 (1984)]. The myriad of known biological activities of IL-1 include the activation of T helper cells, induction of fever, stimulation of prostaglandin or collagenase production, neutrophil chemotaxis, induction of acute phase proteins and the suppression of plasma iron levels.

There are many disease states in which excessive or unregulated IL-1 production is implicated in exacerbating and/or causing the disease. These include rheumatoid arthritis, osteoarthritis, endotoxemia and/or toxic shock syndrome, other acute or chronic inflammatory disease states such as the inflammatory reaction induced by endotoxin or inflammatory bowel disease; tuberculosis, atherosclerosis, muscle degeneration, cachexia, psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, gout, traumatic arthritis, rubella arthritis, and acute synovitis. Recent evidence also links IL-1 activity to diabetes and pancreatic β cells [review of the biological activities which have been attributed to IL-1 Dinarello, *J. Clinical Immunology*, 5 (5), 287-297 (1985)].

Excessive or unregulated TNF production has been implicated in mediating or exacerbating a number of diseases including rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions; sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, adult respiratory distress syndrome, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcoisosis, bone resorption diseases, reperfusion injury, graft vs. host reaction, allograft rejections, fever and myalgias due to infection, such as influenza, cachexia secondary to infection or malignancy, cachexia, secondary to acquired immune deficiency syndrome (AIDS), AIDS, ARC (AIDS related complex), keloid formation, scar tissue formation, Crohn's disease, ulcerative colitis, or pyresis.

Interleukin-8 (IL-8) is a chemotactic factor produced by several cell types including mononuclear cells, fibroblasts, endothelial cells, and keratinocytes. Its production from endothelial cells is induced by IL-1, TNF, or lipopolysaccharide (LPS). IL-8 stimulates a number of functions in vitro. It has been shown to have

chemoattractant properties for neutrophils, T-lymphocytes, and basophils. In addition it induces histamine release from basophils from both normal and atopic individuals as well as lysozomal enzyme release and respiratory burst from neutrophils. IL-8 has also been shown to increase the surface expression of Mac-1 (CD11b/CD18) on neutrophils
5 without de novo protein synthesis, this may contribute to increased adhesion of the neutrophils to vascular endothelial cells. Many diseases are characterized by massive neutrophil infiltration. Conditions associated with an increased in IL-8 production (which is responsible for chemotaxis of neutrophil into the inflammatory site) would benefit by compounds which are suppressive of IL-8 production.

10 IL-1 and TNF affect a wide variety of cells and tissues and these cytokines as well as other leukocyte derived cytokines are important and critical inflammatory mediators of a wide variety of disease states and conditions. The inhibition of these cytokines is of benefit in controlling, reducing and alleviating many of these disease states.

15 Inhibition of signal transduction via CSBP/p38, which in addition to IL-1, TNF and IL-8 described above is also required for the synthesis and/or action of several additional pro-inflammatory proteins (i.e., IL-6, GM-CSF, COX-2, collagenase and stromelysin), is expected to be a highly effective mechanism for regulating the excessive and destructive activation of the immune system. This expectation is
20 supported by the potent and diverse anti-inflammatory activities described for CSBP/p38 kinase inhibitors [Badger, *et al.*, J. Pharm. Exp. Thera. **279** (3): 1453-1461.(1996); Griswold, *et al.*, Pharmacol. Comm. **7**, 323-229 (1996)].

There remains a need for treatment, in this field, for compounds which are cytokine suppressive anti-inflammatory drugs, i.e. compounds which are capable of
25 inhibiting the CSBP/p38/RK kinase.

SUMMARY OF THE INVENTION

This invention relates to the novel compounds of Formula (I) and pharmaceutical compositions comprising a compound of Formula (I) and a
30 pharmaceutically acceptable diluent or carrier.

This invention relates to a method of treating a CSBP/RK/p38 kinase mediated disease in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of Formula (I).

This invention also relates to a method of inhibiting cytokines and the
35 treatment of a cytokine mediated disease, in a mammal in need thereof, which


comprises administering to said mammal an effective amount of a compound of Formula (I).

This invention more specifically relates to a method of inhibiting the production of IL-1 in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I).

This invention more specifically relates to a method of inhibiting the production of IL-8 in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I).

This invention more specifically relates to a method of inhibiting the production of TNF in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I).

Accordingly, the present invention provides a compound of Formula (I):



wherein

R₁ is 4-pyridyl, pyrimidinyl, 4-pyridazinyl, 1,2,4-triazin-5-yl, quinolyl, isoquinolyl, or quinazolin-4-yl ring, which ring is substituted with Y-R_a and optionally with an additional independent substituent selected from C₁₋₄ alkyl, halogen, hydroxyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, C₁₋₄ alkylsulfinyl, CH₂OR₁₂, amino, mono and di- C₁₋₆ alkyl substituted amino, an N-heterocyclyl ring which ring has from 5 to 7 members and optionally contains an additional heteroatom selected from oxygen, sulfur or NR₁₅, N(R₁₀)C(O)R_b or NHR_a;

Y is oxygen or sulfur;

R₄ is phenyl, naphth-1-yl or naphth-2-yl, or a heteroaryl, which is optionally substituted by one or two substituents, each of which is independently selected, and which, for a 4-phenyl, 4-naphth-1-yl, 5-naphth-2-yl or 6-naphth-2-yl substituent, is halogen, cyano, nitro, C(Z)NR₇R₁₇, C(Z)OR₁₆, (CR₁₀R₂₀)_vCOR₁₂, SR₅, SOR₅, OR₁₂, halo-substituted-C₁₋₄ alkyl, C₁₋₄ alkyl, ZC(Z)R₁₂, NR₁₀C(Z)R₁₆, or (CR₁₀R₂₀)_vNR₁₀R₂₀ and which, for other positions of substitution, is halogen, cyano, C(Z)NR₁₃R₁₄, C(Z)OR₃, (CR₁₀R₂₀)_m"COR₃, S(O)_mR₃, OR₃, halo-substituted-C₁₋₄ alkyl, C₁₋₄ alkyl, (CR₁₀R₂₀)_m"NR₁₀C(Z)R₃, NR₁₀S(O)_mR₈, NR₁₀S(O)_m"NR₇R₁₇, ZC(Z)R₃ or (CR₁₀R₂₀)_m"NR₁₃R₁₄;

Z is oxygen or sulfur;

- n is an integer having a value of 1 to 10;
 m is 0, or the integer 1 or 2;
 m' is an integer having a value of 1 or 2,
 m" is 0, or an integer having a value of 1 to 5;
 5 v is 0, or an integer having a value of 1 or 2;
 R₂ is -C(H)(A)(R₂₂);
 A is an optionally substituted aryl, heterocyclyl, or heteroaryl ring, or A is a substituted C₁₋₁₀ alkyl;
 R₂₂ is an optionally substituted C₁₋₁₀ alkyl;
 10 R_a is aryl, arylC₁₋₆alkyl, heterocyclic, heterocyclylC₁₋₆ alkyl, heteroaryl, or heteroarylC₁₋₆alkyl, wherein each of these moieties may be optionally substituted;
 R_b is hydrogen, C₁₋₆ alkyl, C₃₋₇ cycloalkyl, aryl, arylC₁₋₄ alkyl, heteroaryl, heteroarylC₁₋₄alkyl, heterocyclyl, or heterocyclylC₁₋₄ alkyl; and wherein each
 15 of these moieties may be optionally substituted;
 R₃ is heterocyclyl, heterocyclylC₁₋₁₀ alkyl or R₈;
 R₅ is hydrogen, C₁₋₄ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl or NR₇R₁₇, excluding the moieties SR₅ being SNR₇R₁₇ and SOR₅ being SOH;
 R₇ and R₁₇ is each independently selected from hydrogen or C₁₋₄ alkyl or R₇ and
 20 R₁₇ together with the nitrogen to which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR₁₅;
 R₈ is C₁₋₁₀ alkyl, halo-substituted C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₃₋₇ cycloalkyl, C₅₋₇ cycloalkenyl, aryl, arylC₁₋₁₀ alkyl, heteroaryl, heteroarylC₁₋₁₀
 25 alkyl, (CR₁₀R₂₀)_nOR₁₁, (CR₁₀R₂₀)_nS(O)_mR₁₈, (CR₁₀R₂₀)_nNHS(O)₂R₁₈, (CR₁₀R₂₀)_nNR₁₃R₁₄; and wherein the aryl, arylalkyl, heteroaryl, heteroaryl alkyl may be optionally substituted;
 R₉ is hydrogen, C(Z)R₁₁ or optionally substituted C₁₋₁₀ alkyl, S(O)₂R₁₈, optionally substituted aryl or optionally substituted aryl-C₁₋₄ alkyl;
 30 R₁₀ and R₂₀ are each independently selected from hydrogen or C₁₋₄ alkyl;
 R₁₁ is hydrogen, C₁₋₁₀ alkyl, C₃₋₇ cycloalkyl, heterocyclyl, heterocyclyl C₁₋₁₀alkyl, aryl, arylC₁₋₁₀ alkyl, heteroaryl or heteroarylC₁₋₁₀ alkyl, wherein these moieties may be optionally substituted;
 R₁₂ is hydrogen or R₁₆;
 35 R₁₃ and R₁₄ is each independently selected from hydrogen or optionally substituted C₁₋₄ alkyl, optionally substituted aryl or optionally substituted aryl-C₁₋₄ alkyl,

or together with the nitrogen which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR₉ ;

R₁₅ is R₁₀ or C(Z)-C₁₋₄ alkyl;

5 R₁₆ is C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkyl, or C₃₋₇ cycloalkyl;

R₁₈ is C₁₋₁₀ alkyl, C₃₋₇ cycloalkyl, heterocyclyl, aryl, aryl₁₋₁₀alkyl, heterocyclyl, heterocyclyl-C₁₋₁₀alkyl, heteroaryl or heteroaryl₁₋₁₀alkyl;

or a pharmaceutically acceptable salt thereof.

10 DETAILED DESCRIPTION OF THE INVENTION

In Formula (I), suitable R₁ moieties includes 4-pyridyl, 4-pyrimidinyl, 4-pyridazinyl, 1,2,4-triazin-5-yl, 4-quinolyl, 6-isoquinolynyl, or 4-quinazolinyl rings, of which the 4-pyridyl, 4-pyrimidinyl and 4-quinolyl rings are preferred. More preferred is the 4-pyrimidinyl or 4-pyridyl moiety, and most preferred is the 4-

15 pyrimidinyl ring.

The R₁ moiety is substituted with the Y-R_a group wherein Y is oxygen or sulfur, and R_a is aryl, arylC₁₋₆alkyl, heterocyclic, heterocyclicC₁₋₆ alkyl, heteroaryl, or heteroarylC₁₋₆alkyl; and wherein each of the R_a moieties may be optionally substituted as defined below.

20 When R_a is aryl, it is preferably phenyl or naphthyl. When R_a is arylalkyl, it is preferably benzyl or naphthylmethyl. When R_a is a heterocyclic or heterocyclic alkyl moiety, the heterocyclic portion is preferably pyrrolidinyl, piperidinyl, morpholino, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrothiopyransulfinyl, tetrahydrothio-pyransulfonyl, pyrrolidinyl, indole, or piperonyl ring. It is noted

25 that the heterocyclic rings herein may contain unsaturation, such as in a tryptamine ring.

When R_a is a heteroaryl ring as defined below, it is preferably a pyridine or tetrazole ring.

The R_a aryl, heterocyclic and heteroaryl rings may be optionally substituted

30 one or more times, preferably one to three times, independently with halogen; C₁₋₄ alkyl, such as methyl, ethyl, propyl, isopropyl, or t-butyl; halosubstituted alkyl, such as CF₃; hydroxy; hydroxy substituted C₁₋₄ alkyl; (CR₁₀R₂₀)_q C₁₋₄ alkoxy, such as methoxy or ethoxy ; (CR₁₀R₂₀)_q S(O)_malkyl and ; (CR₁₀R₂₀)_qS(O)_m aryl (wherein m is 0, 1, or 2); (CR₁₀R₂₀)_qC(O)OR₁₁, such as C(O)C₁₋₄ alkyl or

35 C(O)OH moieties; (CR₁₀R₂₀)_qC(O)R₁₁; (CR₁₀R₂₀)_qOC(O)R_c; O-(CH₂)_s-O; (CR₁₀R₂₀)_qNR₁₃R₁₄; (CR₁₀R₂₀)_qN(R₁₀)C(O)R_b; (CR₁₀R₂₀)_qC(O)NR₁₃R₁₄;

(CR₁₀R₂₀)_qC(O)NR₁₀R_c; (CR₁₀R₂₀)_qS(O)₂NR₁₃R₁₄;

(CR₁₀R₂₀)_qS(O)₂NR₁₀R_c; (CR₁₀R₂₀)_qN(R₁₀)S(O)₂R_c; cyano, nitro, an

N-heterocyclyl ring which ring has from 5 to 7 members and optionally contains an additional heteroatom selected from oxygen, sulfur or NR₁₅; aryl, such as phenyl;

- 5 an optionally substituted arylalkyl, such as benzyl or phenethyl; aryloxy, such as phenoxy; or arylalkyloxy such as benzyloxy; and wherein the aryl, alkylalkyl, aryloxy and arylalkyloxy containing moieties may be optionally substituted themselves one to two times by halogen, hydroxy, hydroxy substituted alkyl, C₁₋₁₀ alkoxy, S(O)_m alkyl, amino, NR₇R₁₇ group, C₁₋₄ alkyl, or halosubstituted C₁₋₄ alkyl.

Suitably, s is an integer having a value of 1, 2, or 3. Preferably s is 2 yielding a 1,3-dioxyethylene moiety, or ketal functionality.

Suitably, q is 0 or an integer having a value of 1 to 4.

- 15 Suitably, R_b is hydrogen, C₁₋₆ alkyl, C₃₋₇ cycloalkyl, aryl, arylC₁₋₄ alkyl, heteroaryl, heteroarylC₁₋₄alkyl, heterocyclyl, or heterocyclylC₁₋₄ alkyl moiety; all of which moieties may be optionally substituted as defined below.

Suitably, R_c is an C₁₋₆ alkyl, C₃₋₇ cycloalkyl, aryl, arylC₁₋₄ alkyl, heteroaryl, heteroarylC₁₋₄alkyl, heterocyclyl, or heterocyclylC₁₋₄ alkyl moiety, all of which moieties may be optionally substituted as defined below.

- 20 Suitable R_a groups include, but are not limited to, benzyl, halosubstituted benzyl, naphthylmethyl, phenyl, halosubstituted phenyl, aminocarbonylphenyl, alkylphenyl, cyanophenyl, alkylthiophenyl, hydroxyphenyl, alkoxyphenyl, phenoxyphenyl, benzyloxyphenyl, phenylphenyl, methylenedioxyphenyl, trifluoromethylphenyl, methylsulfonylphenyl, tetrazole, methyltetrazolyl,
- 25 morpholinopropyl, piperonyl, piperidin-4-yl, alkyl substituted piperidine, such as 1-methyl piperidine, or 2,2,6,6-tetramethylpiperidin-4-yl.

- Preferred ring substitution on the benzyl or phenyl rings is in the 4-position. Preferred substitution on the phenyl or phenyl alkyl groups is halogen,
- 30 halosubstituted alkyl or alkyl groups, such as fluoro or chloro, or methyl.

- It is recognized that the R₁ group may additionally be substituted one or more times independently by C₁₋₄ alkyl, halo, OH, C₁₋₄ alkoxy, C₁₋₄ alkylthio, C₁₋₄ alkylsulfinyl, CH₂OR₁₂, amino, mono- and di-C₁₋₆ alkyl substituted amino,
- 35 N(R₁₀)C(O)R_b, NHR_a or an N-heterocyclyl ring which ring has from 5 to 7

members and optionally contains an additional heteroatom selected from oxygen, sulfur or NR₁₅.

When the additional R₁ optional substituent is N(R₁₀)C(O)R_b, R_b is preferably C₁₋₆ alkyl; preferably R₁₀ is hydrogen. It is also recognized that the R_b moieties, in particular the C₁₋₆ alkyl group may be optionally substituted, preferably from one to three times, preferably with halogen, such as fluorine, as in trifluoro-methyl or trifluoroethyl.

The preferred ring placement for YR_a moiety on the 4-pyridyl derivative is in the 2-position, and a preferred ring placement on the 4-pyrimidinyl ring is also at the 2-position.

Suitably, R₄ is phenyl, naphth-1-yl or naphth-2-yl, or a heteroaryl, which is optionally substituted by one or two substituents. More preferably R₄ is a phenyl or naphthyl ring. Suitable substitutions for R₄ when this is a 4-phenyl, 4-naphth-1-yl, 5-naphth-2-yl or 6-naphth-2-yl moiety are one or two substituents each of which are independently selected from halogen, SR₅, SOR₅, OR₁₂, CF₃, or (CR₁₀R₂₀)_vNR₁₀R₂₀, and for other positions of substitution on these rings preferred substitution is halogen, S(O)_mR₃, OR₃, CF₃, (CR₁₀R₂₀)_mNR₁₃R₁₄, NR₁₀C(Z)R₃ and NR₁₀S(O)_mR₈.

When R₄ is a heteroaryl ring, the ring is substituted in a similar ring substitution pattern as for the phenyl ring, i.e. halogen, SR₅, SOR₅, OR₁₂, CF₃, or (CR₁₀R₂₀)_vNR₁₀R₂₀.

Preferred substituents for the 4-position in phenyl and naphth-1-yl and on the 5-position in naphth-2-yl include halogen, especially fluoro and chloro and SR₅ and SOR₅ wherein R₅ is preferably a C₁₋₂ alkyl, more preferably methyl; of which the fluoro and chloro is more preferred, and most especially preferred is fluoro.

Preferred substituents for the 3-position in phenyl and naphth-1-yl rings include: halogen, especially fluoro and chloro; OR₃, especially C₁₋₄ alkoxy; CF₃, NR₁₀R₂₀, such as amino; NR₁₀C(Z)R₃, especially NHCO(C₁₋₁₀ alkyl); NR₁₀S(O)_mR₈, especially NHSO₂(C₁₋₁₀ alkyl), and SR₃ and SOR₃ wherein R₃ is preferably a C₁₋₂ alkyl, more preferably methyl. When the phenyl ring is disubstituted preferably it is two independent halogen moieties, such as fluoro and chloro, preferably di-chloro and more preferably in the 3,4-position. It is also preferred that for the 3-position of both the OR₃ and ZC(Z)R₃ moieties, R₃ may also include hydrogen.

Preferably, the R₄ moiety is an unsubstituted or substituted phenyl moiety. More preferably, R₄ is phenyl or phenyl substituted at the 4-position with fluoro

and/or substituted at the 3-position with fluoro, chloro, C₁₋₄ alkoxy, methane-sulfonamido or acetamido, or R₄ is a phenyl di-substituted at the 3,4-position independently with chloro or fluoro, more preferably chloro. Most preferably, R₄ is a 4-fluorophenyl.

- 5 Suitably, Z is oxygen or sulfur, preferably oxygen.
 Suitably, R₃ is heterocyclyl, heterocyclylC₁₋₁₀ alkyl or R₈.
 Suitably, R₅ is hydrogen, C₁₋₄ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl or NR₇R₁₇, excluding the moieties SR₅ being SNR₇R₁₇ and SOR₅ being SOH.
 Suitably, R₆ is hydrogen, a pharmaceutically acceptable cation, C₁₋₁₀ alkyl, C₃₋₇ cycloalkyl, aryl, arylC₁₋₄ alkyl, heteroaryl, heteroarylC₁₋₄alkyl, heterocyclyl, aroyl, or C₁₋₁₀ alkanoyl.
- 10 Suitably, R₇ and R₁₇ is each independently selected from hydrogen or C₁₋₄ alkyl or R₇ and R₁₇ together with the nitrogen to which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR₁₅.
- 15 Suitably, R₈ is C₁₋₁₀ alkyl, halo-substituted C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₃₋₇ cycloalkyl, C₅₋₇ cycloalkenyl, aryl, arylC₁₋₁₀ alkyl, heteroaryl, heteroarylC₁₋₁₀ alkyl, (CR₁₀R₂₀)_nOR₁₁, (CR₁₀R₂₀)_nS(O)_mR₁₈, (CR₁₀R₂₀)_nNHS(O)₂R₁₈, (CR₁₀R₂₀)_nNR₁₃R₁₄; wherein the aryl, arylalkyl, heteroaryl, and heteroarylalkyl containing moieties may be optionally substituted.
- 20 Suitably, R₉ is hydrogen, C(Z)R₁₁ or optionally substituted C₁₋₁₀ alkyl, S(O)₂R₁₈, optionally substituted aryl or an optionally substituted aryl-C₁₋₄ alkyl.
 Suitably, R₁₀ and R₂₀ are each independently selected from hydrogen or C₁₋₄ alkyl.
- 25 Suitably, R₁₁ is hydrogen, C₁₋₁₀ alkyl, C₃₋₇ cycloalkyl, heterocyclyl, heterocyclyl C₁₋₁₀alkyl, aryl, arylC₁₋₁₀ alkyl, heteroaryl or heteroarylC₁₋₁₀ alkyl; and wherein all of these moieties may be optionally substituted.
 Suitably, R₁₂ is hydrogen or R₁₆; and R₁₆ is suitably, C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkyl, or C₃₋₇ cycloalkyl.
- 30 Suitably, R₁₃ and R₁₄ is each independently selected from hydrogen or optionally substituted C₁₋₄ alkyl, optionally substituted aryl or optionally substituted aryl-C₁₋₄ alkyl, or together with the nitrogen which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR₉.
- 35 Suitably, R₁₅ is R₁₀ or C(Z)-C₁₋₄ alkyl.

Suitably, R₁₈ is C₁₋₁₀ alkyl, C₃₋₇ cycloalkyl, heterocyclyl, aryl, aryl₁₋₁₀alkyl, heterocyclyl, heterocyclyl-C₁₋₁₀alkyl, heteroaryl or heteroaryl₁₋₁₀alkyl.

Suitably, v is 0, or an integer having a value of 1 or 2.

5 Suitably, m is 0, or the integer 1 or 2.

Suitably, m' is an integer having a value of 1 or 2.

Suitably, m" is 0, or an integer having a value of 1 to 5.

Suitably, n is an integer having a value of 1 to 10.

10 R₂ is a substituted alkyl derivative. It is recognized that the first methylene carbon in this chain is a tertiary carbon, and it will contain one hydrogen moiety. This methylene group will have has two additional substituents, an R₂₂ moiety and an A moiety, i.e., C(H)(A)(R₂₂). Both A and R₂₂ may not be unsubstituted C₁₋₁₀ alkyl moieties.

15 In a preferred embodiment, R₂ is a C(AA₁)(A) moiety, wherein AA₁ is the R₂₂ moiety, but is specifically the side chain residue (R) of an amino acid, as is further described herein.

Suitably, A is an optionally substituted C₃₋₇cycloalkyl, aryl, heteroaryl, or heterocyclic ring, or A is a substituted C₁₋₁₀ alkyl moiety.

20 When A is an aryl, heteroaryl and heterocyclic ring, the ring may be substituted independently one or more times, preferably, 1 to 3 times by C₁₋₁₀ alkyl; halogen; halo substituted C₁₋₁₀ alkyl, such as CF₃; (CR₁₀R₂₀)_tOR₁₁; (CR₁₀R₂₀)_tNR₁₃R₁₄, especially amino or mono- or di-C₁₋₄ alkylamino; (CR₁₀R₂₀)_tS(O)_mR₁₈, wherein m is 0, 1 or 2; SH; NR₁₀C(Z)R₃ (such NHCO(C₁₋₁₀ alkyl)); or NR₁₀S(O)_mR₈ (such as
25 NHSO₂(C₁₋₁₀ alkyl)).

Suitably, t is 0, or an integer of 1 to 4.

When A is an optionally substituted cycloalkyl it is as defined below in the R₂₂ substitution.

30 When A is an optionally substituted heterocyclyl ring, the ring is preferably a morpholino, pyrrolidinyl, piperazinyl or a piperidinyl ring.

When A is an optionally substituted aryl moiety, it is preferably a phenyl ring.

When A is an optionally substituted heteroaryl ring, the heteroaryl term is as defined below in the definition section.

35 When A is a substituted C₁₋₁₀ alkyl moiety, the alkyl chain may be straight or branched. The chain is substituted independently 1 or more times, preferably 1 to 3 times by halogen, such as fluorine, chlorine, bromine or iodine; halosubstituted C₁₋₁₀

alkyl, such as CF₃; C₃₋₇cycloalkyl, C₁₋₁₀ alkoxy, such as methoxy or ethoxy; hydroxy substituted C₁₋₁₀ alkoxy; halosubstituted C₁₋₁₀ alkoxy, such as OCF₂CF₂H; OR₁₁; S(O)_mR₁₈ (wherein m is 0, 1 or 2); NR₁₃R₁₄; C(Z)NR₁₃R₁₄; S(O)_mNR₁₃R₁₄; NR₂₃C(Z)R₁₁; NHS(O)₂R₁₈; C(Z)R₁₁; OC(Z)R₁₁; C(Z)OR₁₁; 5 C(Z)NR₁₁OR₉; N(OR₆)C(Z)NR₁₃R₁₄; N(OR₆)C(Z)R₁₁; C(=NOR₆)R₁₁; NR₂₃C(=NR₁₉)NR₁₃R₁₄; OC(Z)NR₁₃R₁₄; NR₂₃C(Z)NR₁₃R₁₄; or NR₂₃C(Z)OR₁₀.

Preferably, A is a C₃₋₇ cycloalkyl, or a C₁₋₆ alkyl, more preferably a C₁₋₂ alkyl, i.e. a methylene or ethylene moiety, more preferably a methylene moiety which 10 is substituted by one of the above noted groups.

Preferably, when A is an alkyl derivative, it is substituted by OR₁₁ where R₁₁ is preferably hydrogen, aryl or arylalkyl; NR₁₃R₁₄; OC(Z)R₁₁; or C(Z)OR₁₁.

More preferably, A is substituted by OR₁₁ where R₁₁ is hydrogen.

Suitably, R₂₂ is a C₁₋₁₀ alkyl chain, which chain may be straight or branched 15 and which may be optionally substituted independently, one or more times, preferably 1 to 3 times, by halogen, such as fluorine, chlorine, bromine or iodine; halo substituted C₁₋₁₀ alkyl; C₁₋₁₀ alkoxy, such as methoxy or ethoxy; hydroxy substituted C₁₋₁₀ alkoxy; halosubstituted C₁₋₁₀ alkoxy, such as OCF₂CF₂H; OR₁₁; S(O)_mR₁₈; NR₁₃R₁₄; C(Z)NR₁₃R₁₄; S(O)_mNR₁₃R₁₄; NR₂₃C(Z)R₁₁; NHS(O)₂R₁₈; 20 C(Z)R₁₁; OC(Z)R₁₁; C(Z)OR₁₁; C(Z)NR₁₁OR₉; N(OR₆)C(Z)NR₁₃R₁₄; N(OR₆)C(Z)R₁₁; C(=NOR₆)R₁₁; NR₂₃C(=NR₁₉)NR₁₃R₁₄; OC(Z)NR₁₃R₁₄; NR₂₃C(Z)NR₁₃R₁₄; NR₂₃C(Z)OR₁₀; optionally substituted C₃₋₇ cycloalkyl; optionally substituted aryl, such as phenyl; optionally substituted heteroaryl; or an optionally substituted heterocyclic moiety. The optional substituents on these 25 cycloalkyl, aryl, heteroaryl, and heterocyclic moieties are as defined herein below.

It is noted that those R₂₂ substituent groups which contain carbon as the first connecting group, i.e. C(Z)OR₁₁; C(Z)NR₁₁OR₉; C(Z)R₁₁; C(Z)NR₁₃R₁₄; C(=NOR₆)R₁₁, may be the sole carbon in alkyl chain. Therefore, R₂₂ may, for instance, be a carboxy, an aldehyde, an amide, as well as being a substituent off a 30 methylene unit, such as carbamoylmethyl, or acetamidomethyl. In other words, R₂₂ can be an optionally substituted alkyl group as defined above, or R₂₂ can be C(Z)OR₁₁, C(Z)NR₁₁OR₉, C(Z)R₁₁, C(Z)NR₁₃R₁₄, or C(=NOR₆)R₁₁.

Preferably R₂₂ is a C₁₋₆ unsubstituted or substituted alkyl group, such as a C₁₋₃ alkylene, such as methyl, ethyl or isopropyl, or a methylene or ethylene moiety 35 substituted by one of the above noted moieties, or as noted above those substituent groups which contain a carbon may substituent for the first methylene unit of the alkyl

chain, such as carboxy, $C(O)OR_{11}$, $C(O)NR_{13}R_{14}$, or R_{22} is an optionally substituted aryl group, such as a benzyl or phenethyl.

Preferably R_{22} is a C_{1-6} unsubstituted or substituted alkyl group, more preferably a C_{1-2} alkylene chain, such as a methylene or ethylene moiety, more preferably methylene.

Preferably the R_{22} alkyl chain is substituted by OR_{11} , where R_{11} is preferably hydrogen, aryl or arylalkyl; $S(O)mR_{18}$, where m is 0 and R_{18} is a C_{1-6} alkyl; or an optionally substituted aryl, i.e. a benzyl or phenethyl moiety.

More preferably, R_{22} is methyl, phenyl, benzyl, CH_2OH , or CH_2-O -aryl.

Preferably, one or both of A and R_{22} contain hydroxy moieties, such as in C_{1-6} alkyl OR_{11} , wherein R_{11} is hydrogen, i.e. CH_2CH_2OH .

Suitably, when AA_1 is the (R) side chain residue of an amino acid, it is a C_{1-6} alkyl group, which may be straight or branched. This means the R group off the core amino acid of the structure $R-C(H)(COOH)(NH_2)$. The R residue term is for example, CH_3 for alanine, $(CH_3)_2CH-$ for valine, $(CH_3)_2CH-CH_2-$ for leucine, phenyl- CH_2- for phenylalanine, $CH_3-S-CH_2-CH_2-$ for methionine, etc. All generally recognized primary amino acids are included in this groups, such as but not limited to, alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, serine, threonine, tryptophan, tyrosine, valine, hydroxylysine, methylhistidine, and other naturally occurring amino acids not found in proteins, such as β -alanine, γ -aminobutyric acid, homocysteine, homoserine, citrulline, ornithine, canavanine, djenkolic acid, and β -cyanoalanine, or other naturally occurring non-mammalian amino acids.

Preferably AA_1 is the residue of phenylalanine, or alanine.

When R_{22} is an optionally substituted heterocyclic moiety, the ring is preferably a morpholino, pyrrolidinyl, piperazinyl, or a piperidinyl group. When the heterocyclic ring is optionally substituted the substituents may be directly attached to the free nitrogen, such as in the piperidinyl group or pyrrole ring, or on the ring itself. Preferably the ring is a piperidine or pyrrole, more preferably piperidine.

The R_{22} heterocyclyl ring may be optionally substituted one to four times independently by halogen; C_{1-4} alkyl; aryl, such as phenyl; arylalkyl, such as benzyl, (and wherein the aryl or aryl alkyl moieties themselves may be optionally substituted as defined in the definition section below); $C(O)OR_{11}$, such as the $C(O)C_{1-4}$ alkyl or $C(O)OH$ moieties; $C(O)H$; $C(O)C_{1-4}$ alkyl; hydroxy substituted C_{1-4} alkyl; C_{1-4} alkoxy; $S(O)_mC_{1-4}$ alkyl (wherein m is 0, 1, or 2); or $NR_{10}R_{20}$ (wherein R_{10} and R_{20} are independently hydrogen or C_{1-4} alkyl).

Preferably if the ring is a piperidine, the substituents are attached directly on the available nitrogen, i.e. a 1-Formyl-4-piperidine, 1-benzyl-4-piperidine, 1-methyl-4-piperidine, 1-ethoxycarbonyl-4-piperidine. If the ring is substituted by an alkyl group and the ring is attached in the 4-position, it is preferably substituted in the 2- or 6-
 5 position or both, such as 2,2,6,6-tetramethyl-4-piperidine. Similarly, if the ring is a pyrrole, the substituents are all directly on the available nitrogen.

When the R₂₂ optional substituent is an optionally substituted aryl, it is preferably a phenyl; or when R₂₂ is an optionally substituted heteroaryl ring (as defined in the definition section below), the rings may be optionally substituted
 10 independently one or more times, preferably by one to three times by C₁₋₁₀ alkyl; halogen, especially fluoro or chloro; (CR₁₀R₂₀)_tOR₁₁; (CR₁₀R₂₀)_tNR₁₃R₁₄; especially amino or mono- or di-C₁₋₄ alkylamino; (CR₁₀R₂₀)_tS(O)_mR₁₈, wherein m is 0, 1 or 2; SH; OR₁₁; NR₁₀C(Z)R₃ (such as NHCO(C₁₋₁₀ alkyl)); or NR₁₀S(O)_mR₈ (such as NHSO₂(C₁₋₁₀ alkyl)).

When A or R₂₂ is an (optionally) substituted C₃₋₇cycloalkyl group, it is preferably a C₃ or C₆ ring, most preferably a C₃ ring, which ring may be optionally substituted one or more time, preferably 1 to 3 times, independently by halogen, such as fluorine, or chlorine; (CR₁₀R₂₀)_tOR₁₁; S(O)_mR₁₈; cyano, (CR₁₀R₂₀)_tNR₁₃R₁₄, especially amino or mono- or di-C₁₋₄ alkylamino; N(R₁₀)C(O)X₁ and X₁ is C₁₋₄
 20 alkyl, aryl or arylC₁₋₄alkyl; C₁₋₁₀ alkyl, such as methyl, ethyl, propyl, isopropyl, or t-butyl; an optionally substituted alkyl wherein the substituents are halogen, (such as CF₃), hydroxy, nitro, cyano, amino, NR₁₃R₁₄, or S(O)_mR₁₈; an optionally substituted alkylene, such as ethylene or propylene; an optionally substituted alkyne, such as ethyne; C(O)OR₁₁; the group R_e; C(O)H; =O; =N-OR₁₁; N(H)-OH (or
 25 substituted alkyl or aryl derivatives thereof on the nitrogen or the oxime moiety); or N(OR_d)-C(O)-R_f.

Suitably R_d is hydrogen, a pharmaceutically acceptable cation, aroyl or a C₁₋₁₀ alkanoyl group.

Suitably R_e is a 1,3-dioxyalkylene group of the formula -O-(CH₂)_s-O-, wherein
 30 s is 1 to 3, preferably s is 2 yielding a 1,3-dioxyethylene moiety, or ketal functionality.

Suitably R_f is NR₂₁R₂₄; alkyl 1-6; halosubstituted alkyl 1-6; hydroxy substituted alkyl 1-6; alkenyl 2-6; aryl or heteroaryl optionally substituted by halogen, alkyl 1-6, halosubstituted alkyl 1-6, hydroxyl, or alkoxy 1-6.

Suitably R₂₁ is hydrogen, or alkyl 1-6.

35 Suitably R₂₄ is hydrogen, alkyl 1-6, aryl, benzyl, heteroaryl, alkyl substituted by halogen or hydroxyl, or phenyl substituted by a member selected from the group

consisting of halo, cyano, alkyl₁₋₆, alkoxy₁₋₆, halosubstituted alkyl₁₋₆, S(O)_m alkyl₁₋₆; or R₂₁ and R₂₄ may together with the nitrogen to which they are attached form a ring having 5 to 7 members, which members may be optionally replaced by a heteroatom selected from oxygen, sulfur or nitrogen. The ring may be saturated or contain more than one unsaturated bond. Preferably R_f is NR₂₁R₂₄, and more preferably R₂₁ and R₂₄ are both hydrogen.

When the A or R₂₂ optional substituent is NR₁₃R₁₄ it is recognized that in some instances this can yield the same moiety as a heterocyclic moiety noted above which is also a suitable variable. Preferably R₁₃ and R₁₄ are independently hydrogen, C₁₋₄ alkyl, preferably methyl, or benzyl.

When the A or R₂₂ optional substituent is a C(Z)OR₁₁ group, R₁₁ is suitably hydrogen, C₁₋₄ alkyl, especially methyl.

When the A or R₂₂ optional substituent is a S(O)_mR₁₈ group, R₁₈ is preferably aryl, especially phenyl, or a C₁₋₁₀ alkyl, especially methyl, or ethyl.

When the A or R₂₂ optional substituent is a OR₁₁ group, R₁₁ is preferably hydrogen, aryl, especially phenyl, or C₁₋₁₀ alkyl, especially methyl or ethyl.

When the A or R₂₂ optional substituent is a NHS(O)₂R₁₈ group, R₁₈ is suitably alkyl, especially methyl.

In a preferred subgenus of compounds of Formula (I), R₁ is 2-phenoxy-4-pyridyl or 2-phenoxy-4-pyrimidinyl. R₄ is phenyl or phenyl substituted one or two times by fluoro, chloro, C₁₋₄ alkoxy, -S(O)_m alkyl, methanesulfonamido or acetamido. A is CH₂OH, CH₃, benzyl, phenyl, CH₂NH(methyl) or CH₂N(dimethyl); and R₂₂ is methyl, ethyl, methanol, or CH₂-O-phenyl; or alternatively the R₂ group is 1-hydroxy-3-phenylprop-2-yl, 1-hydroxyprop-2-yl, 1-hydroxybut-2-yl, 1,3-dihydroxyprop-2-yl, or 1-hydroxy-2-phenylethyl-2-yl.

As used herein, "optionally substituted" unless specifically defined shall mean such groups as halogen, such as fluorine, chlorine, bromine or iodine; hydroxy; hydroxy substituted C₁₋₁₀alkyl; C₁₋₁₀ alkoxy, such as methoxy or ethoxy; halosubstituted C₁₋₁₀ alkoxy; S(O)_m alkyl, such as methyl thio, methylsulfinyl or methyl sulfonyl; NR₇R₁₇, such as amino or mono or -disubstituted C₁₋₄ alkyl or wherein the R₇R₁₇ can cyclize together with the nitrogen to which they are attached to form a 5 to 7 membered ring which optionally contains an additional heteroatom selected from O/N/S; C₁₋₁₀ alkyl, C₃₋₇cycloalkyl, or C₃₋₇cycloalkyl C₁₋₁₀ alkyl group, such as methyl, ethyl, propyl, isopropyl, t-butyl, etc. or cyclopropyl methyl;

halosubstituted C₁₋₁₀ alkyl, such CF₂CF₂H, or CF₃; an optionally substituted aryl, such as phenyl, or an optionally substituted arylalkyl, such as benzyl or phenethyl, wherein these aryl moieties may also be substituted one to two times by halogen; hydroxy; hydroxy substituted alkyl; C₁₋₁₀ alkoxy; S(O)_malkyl; amino, mono & di-substituted C₁₋₄ alkyl amino, such as in the NR₇R₁₇ group; C₁₋₄ alkyl, or CF₃.

Suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of inorganic and organic acids, such as hydrochloric acid, hydrobromic acid, sulphuric acid, phosphoric acid, methane sulphonic acid, ethane sulphonic acid, acetic acid, malic acid, tartaric acid, citric acid, lactic acid, oxalic acid, succinic acid, fumaric acid, maleic acid, benzoic acid, salicylic acid, phenylacetic acid and mandelic acid.

In addition, pharmaceutically acceptable salts of compounds of Formula (I) may also be formed with a pharmaceutically acceptable cation, for instance, if a substituent group comprises a carboxy moiety. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations.

The term "halo" or "halogens" is used herein to mean the halogens, chloro, fluoro, bromo and iodo.

The term "C₁₋₁₀alkyl" or "alkyl" is used herein to mean both straight and branched chain radicals of 1 to 10 carbon atoms, unless the chain length is otherwise limited, including, but not limited to, methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *tert*-butyl, *n*-pentyl and the like.

The term "cycloalkyl" is used herein to mean cyclic radicals, preferably of 3 to 8 carbons, including but not limited to cyclopropyl, cyclopentyl, cyclohexyl, and the like.

The term "cycloalkenyl" is used herein to mean cyclic radicals, preferably of 5 to 8 carbons, which have at least one bond including but not limited to cyclopentenyl, cyclohexenyl, and the like.

The term "alkenyl" is used herein at all occurrences to mean straight or branched chain radical of 2-10 carbon atoms, unless the chain length is limited thereto, including, but not limited to ethenyl, 1-propenyl, 2-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl and the like.

The term "aryl" is used herein to mean phenyl and naphthyl;

The term "heteroaryl" (on its own or in any combination, such as "heteroaryloxy", or "heteroaryl alkyl") is used herein to mean a 5-10 membered aromatic ring system in which one or more rings contain one or more heteroatoms selected from the group consisting of N, O or S, such as, but not limited, to pyrrole, pyrazole, furan, thiophene, quinoline, isoquinoline, quinazolinyl, pyridine, pyrimidine, oxazole, thiazole, thiadiazole, tetrazole, triazole, imidazole, or benzimidazole.

The term "heterocyclic" (on its own or in any combination, such as "heterocyclalkyl") is used herein to mean a saturated or partially unsaturated 4-10 membered ring system in which one or more rings contain one or more heteroatoms selected from the group consisting of N, O, or S; such as, but not limited to, pyrrolidine, piperidine, piperazine, morpholine, tetrahydro pyran, or imidazolidine.

The term "aralkyl" or "heteroarylalkyl" or "heterocyclicalkyl" is used herein to mean C₁₋₄ alkyl as defined above attached to an aryl, heteroaryl or heterocyclic moiety as also defined herein unless otherwise indicate.

The term "sulfinyl" is used herein to mean the oxide S (O) of the corresponding sulfide, the term "thio" refers to the sulfide, and the term "sulfonyl" refers to the fully oxidized S (O)₂ moiety.

The term "aroyl" is used herein to mean C(O)Ar, wherein Ar is as phenyl, naphthyl, or aryl alkyl derivative such as defined above, such group include but are not limited to benzyl and phenethyl.

The term "alkanoyl" is used herein to mean C(O)C₁₋₁₀ alkyl wherein the alkyl is as defined above.

It is recognized that the compounds of the present invention may exist as stereoisomers, regioisomers, or diastereoisomers. These compounds may contain one or more asymmetric carbon atoms and may exist in racemic and optically active forms. All of these compounds are included within the scope of the present invention.

30

Exemplified compounds of Formula (I) include:

1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-fluorophenoxy)pyrimidin-4-yl]imidazole

35

1-(1,3-Dihydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-fluorophenoxy)pyrimidin-4-yl]imidazole

1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy)pyrimidin-4-yl)imidazole

- 1-(1-Phenoxyprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole
 1-(1-Hydroxy-3-phenylprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole
 1-(1-Hydroxybut-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole
 5 1-(1,3-Dihydroxyprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole
 1-(1-Hydroxy-2-phenyleth-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole
 1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-chlorophenoxy)pyrimidin-4-yl]imidazole
 10 1-(1-Hydroxybut-2-yl)-4-(4-fluorophenyl)-5-[2-(4-chlorophenoxy)pyrimidin-4-yl]imidazole
 1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-methylphenoxy)pyrimidin-4-yl]imidazole
 15 1-(1-Hydroxybut-2-yl)-4-(4-fluorophenyl)-5-[2-(4-methylphenoxy)pyrimidin-4-yl]imidazole
 1-(1,3-Dihydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-methylphenoxy)pyrimidin-4-yl]imidazole
 1-(4-N-Morpholinyl-2-but-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole
 20

The compounds of Formula (I) may be obtained by applying synthetic procedures, some of which are illustrated in Schemes I to XII herein. The synthesis provided for in these Schemes is applicable for the producing compounds of Formula (I) having a variety of different R₁, R₂, and R₄ groups which are reacted, employing optional substituents which are suitably protected, to achieve compatibility with the reactions outlined herein. Subsequent deprotection, in those cases, then affords compounds of the nature generally disclosed. While the schemes describe compounds of Formula (I) with Y as oxygen, one skilled in the art would readily be able to make compounds of Formula (I) wherein Y is sulfur using similar reaction processes as exemplified herein.

Once the imidazole nucleus has been established, further compounds of Formula (I) may be prepared by applying standard techniques for functional group interconversion, well known in the art. For instance: C(O)NR₁₃R₁₄ from CO₂CH₃ by heating with or without catalytic metal cyanide, e.g. NaCN, and HNR₁₃R₁₄ in CH₃OH; OC(O)R₃ from OH with e.g., ClC(O)R₃ in pyridine; NR₁₀-C(S)NR₁₃R₁₄

from NHR_{10} with an alkylisothiocyanate or thiocyanic acid; $\text{NR}_6\text{C(O)OR}_6$ from NHR_6 with the alkyl chloroformate; $\text{NR}_{10}\text{C(O)NR}_{13}\text{R}_{14}$ from NHR_{10} by treatment with an isocyanate, e.g. $\text{HN}=\text{C}=\text{O}$ or $\text{R}_{10}\text{N}=\text{C}=\text{O}$; $\text{NR}_{10}\text{-C(O)R}_8$ from NHR_{10} by treatment with Cl-C(O)R_3 in pyridine; $\text{C(=NR}_{10})\text{NR}_{13}\text{R}_{14}$ from $\text{C(NR}_{13}\text{R}_{14})\text{SR}_3$ with $\text{H}_3\text{NR}_3^+\text{OAc}^-$ by heating in alcohol; $\text{C(NR}_{13}\text{R}_{14})\text{SR}_3$ from $\text{C(S)NR}_{13}\text{R}_{14}$ with $\text{R}_6\text{-I}$ in an inert solvent, e.g. acetone; $\text{C(S)NR}_{13}\text{R}_{14}$ (where R_{13} or R_{14} is not hydrogen) from C(S)NH_2 with $\text{HNR}_{13}\text{R}_{14}\text{-C(=NCN)-NR}_{13}\text{R}_{14}$ from $\text{C(=NR}_{13}\text{R}_{14})\text{-SR}_3$ with NH_2CN by heating in anhydrous alcohol, alternatively from $\text{C(=NH)-NR}_{13}\text{R}_{14}$ by treatment with BrCN and NaOEt in EtOH ;

10 $\text{NR}_{10}\text{-C(=NCN)SR}_8$ from NHR_{10} by treatment with $(\text{R}_8\text{S})_2\text{C=NCN}$; $\text{NR}_{10}\text{SO}_2\text{R}_3$ from NHR_{10} by treatment with ClSO_2R_3 by heating in pyridine; $\text{NR}_{10}\text{C(S)R}_3$ from $\text{-NR}_{10}\text{C(O)R}_8$ by treatment with Lawesson's reagent [2,4-bis(4-methoxyphenyl)-1,3,2,4-dithiadiphosphetane-2,4-disulfide]; $\text{NR}_{10}\text{SO}_2\text{CF}_3$ from NHR_6 with triflic anhydride and base wherein R_3 , R_6 , R_{10} , R_{13} and R_{14} are as defined in Formula (I)


15 herein.

Precursors of the groups R_1 , R_2 and R_4 can be other R_1 , R_2 and R_4 groups which can be interconverted by applying standard techniques for functional group interconversion. For example a compound of the formula (I) wherein R_2 is halo substituted C_{1-10} alkyl can be converted to the corresponding C_{1-10} alkyl N_3 derivative by reacting with a suitable azide salt, and thereafter if desired can be reduced to the corresponding C_{1-10} alkyl NH_2 compound, which in turn can be reacted with $\text{R}_{18}\text{S(O)}_2\text{X}$ wherein X is halo (e.g., chloro) to yield the corresponding C_{1-10} alkyl $\text{NHS(O)}_2\text{R}_{18}$ compound.

20

Alternatively a compound of the formula (I) where R_2 is halo-substituted C_{1-10} -alkyl can be reacted with an amine $\text{R}_{13}\text{R}_{14}\text{NH}$ to yield the corresponding C_{1-10} -alkyl $\text{NR}_{13}\text{R}_{14}$ compound, or can be reacted with an alkali metal salt of R_{18}SH to yield the corresponding C_{1-10} alkyl SR_{18} compound.

25



Scheme I

5


Referring to Scheme I the compounds of Formula (I) are suitably prepared by reacting a compound of the Formula (II) with a compound of the Formula (III) wherein p is 0 or 2, R_1 , R_2 and R_4 are as defined herein, for Formula (I), or are precursors of the groups R_1 , R_2 and R_4 , and Ar is an optionally substituted phenyl group, and thereafter if necessary converting a precursor of R_1 , R_2 and R_4 to a group R_1 , R_2 and R_4 . It is recognized that R_2NH_2 which is reacted with R_1CHO to form

10

the imine, Formula (III) the R₂ moiety when it contains a reactive functional group, such as a primary or secondary amine, an alcohol, or thiol compound the group must be suitably protected. Suitable protecting groups may be found in, Protecting Groups in Organic Synthesis, Greene T W, Wiley-Interscience, New York, 1981, whose disclosure is incorporated herein by reference. For instance, when R₂ is a heterocyclic ring, such as a piperidine ring, the nitrogen is protected with groups such as t-Boc, CO₂R₁₈, or a substituted arylalkyl moiety.

Suitably, the reaction is performed at ambient temperature or with cooling (e.g. -50° to 10°) or heating in an inert solvent such as methylene chloride, DMF, tetrahydrofuran, toluene, acetonitrile, or dimethoxyethane in the presence of an appropriate base such as K₂CO₃, t-buNH₂, 1,8-diazabicyclo [5.4.0.] undec-7-ene (DBU), or a guanidine base such as 1,5,7-triaza-bicyclo [4.4.0] dec-5-ene (TBD). The intermediates of formula (II) have been found to be very stable and capable of storage for a long time. Preferably, p is 2. PTC is defined as a phase transfer catalyst for use herein.

Compounds of the Formula (II) have the structure:




wherein p is 0, or 2; R₄ is as defined for Formula (I) and Ar is an optionally substituted aryl as defined herein. Suitably, Ar is phenyl optionally substituted by C₁₋₄alkyl, C₁₋₄ alkoxy or halo. Preferably Ar is phenyl or 4-methylphenyl, i.e. a tosyl derivative.

Reaction a compound of the Formula (II) wherein p = 2, with a compound of the Formula (III)-Scheme I gives consistently higher yields of compounds of Formula (I) than when p=0. In addition, the reaction of Formula (II) compounds wherein p = 2 is more environmentally and economically attractive. When p=0, the preferred solvent used is methylene chloride, which is environmentally unattractive for large scale processing, and the preferred base, TBD, is also expensive, and produces some byproducts and impurities, than when using the commercially attractive synthesis (p=2) as further described herein.

As noted, Scheme I utilizes the 1,3-dipolar cycloadditions of an anion of a substituted aryl thiomethylisocyanide (when p=0) to an imine. More specifically, this reaction requires a strong base, such as an amine base, to be used for the deprotonation step. The commercially available TBD is preferred although

t-butoxide, Li⁺ or Na⁺, or K⁺ hexamethyldisilazide may also be used. While methylene chloride is the preferred solvent, other halogenated solvents, such as chloroform or carbon tetrachloride; ethers, such as THF, DME, DMF, diethylether, t-butyl methyl ether; as well as acetonitrile, toluene or mixtures thereof can be
5 utilized. The reaction may take place from about -20°C to about; 40°C, preferably from about 0°C to about 23°C, more preferably from about 0°C to about 10°C, and most preferably about 4°C for reactions involving an R₁ group of pyrimidine. For compounds wherein R₁ is pyridine, it is recognized that varying the reactions
10 conditions of both temperature and solvent may be necessary, such as decreasing temperatures to about -50°C or changing the solvent to THF.

In a further process, compounds of Formula (I) may be prepared by coupling a suitable derivative of a compound of Formula (IX):



15 wherein T₁ is hydrogen and T₄ is R₄, or alternatively T₁ is R₁ and T₄ is H in which R₁, R₂ and R₄ are as hereinbefore defined; with: (i) when T₁ is hydrogen, a suitable derivative of the heteroaryl ring R₁H, under ring coupling conditions, to effect coupling of the heteroaryl ring R₁ to the imidazole nucleus at position 5; (ii) when T₄ is
20 hydrogen, a suitable derivative of the aryl ring R₄H, under ring coupling conditions, to effect coupling of the aryl ring R₄ to the imidazole nucleus at position 4.

Such aryl/heteroaryl coupling reactions are well known to those skilled in the art. In general, an organometallic synthetic equivalent of an anion of one component is coupled with a reactive derivative of the second component, in the presence of a
25 suitable catalyst. The anion equivalent may be formed from either the imidazole of Formula (IX), in which case the aryl/heteroaryl compound provides the reactive derivative, or the aryl/heteroaryl compound in which case the imidazole provides the reactive derivative. Accordingly, suitable derivatives of the compound of Formula (IX) or the aryl/heteroaryl rings include organometallic derivatives such as
30 organomagnesium, organozinc, organostannane and boronic acid derivatives and suitable reactive derivatives include the bromo, iodo, fluorosulfonate and trifluoromethanesulphonate derivatives. Suitable procedures are described in WO 91/19497, the disclosure of which is incorporated by reference herein.

Suitable organomagnesium and organozinc derivatives of a compound of Formula (IX) may be reacted with a halogen, fluorosulfonate or triflate derivative of the heteroaryl or aryl ring, in the presence of a ring coupling catalyst, such as a palladium (O) or palladium (II) catalyst, following the procedure of Kumada *et al.*,
5 Tetrahedron Letters, **22**, 5319 (1981). Suitable such catalysts include *tetrakis*-(triphenylphosphine)palladium and PdCl₂[1,4-*bis*-(diphenylphosphino)-butane], optionally in the presence of lithium chloride and a base, such as triethylamine. In addition, a nickel (II) catalyst, such as Ni(II)Cl₂(1,2-biphenylphosphino)ethane, may also be used for coupling an aryl ring, following the procedure of Pridgen *et al.*, J.
10 Org. Chem, 1982, **47**, 4319. Suitable reaction solvents include hexamethylphosphor-amide. When the heteroaryl ring is 4-pyridyl, suitable derivatives include 4-bromo- and 4-iodo-pyridine and the fluorosulfonate and triflate esters of 4-hydroxy pyridine. Similarly, suitable derivatives for when the aryl ring is phenyl include the bromo, fluorosulfonate, triflate and, preferably, the iodo-
15 derivatives. Suitable organomagnesium and organozinc derivatives may be obtained by treating a compound of Formula (IX) or the bromo derivative thereof with an alkyl lithium compound to yield the corresponding lithium reagent by deprotonation or transmetalation, respectively. This lithium intermediate may then be treated with an excess of a magnesium halide or zinc halide to yield the corresponding
20 organometallic reagent.

A trialkyltin derivative of the compound of Formula (IX) may be treated with a bromide, fluorosulfonate, triflate, or, preferably, iodide derivative of an aryl or heteroaryl ring compound, in an inert solvent such as tetrahydrofuran, preferably containing 10% hexamethylphosphoramide, in the presence of a suitable coupling
25 catalyst, such as a palladium (O) catalyst, for instance *tetrakis*-(triphenylphosphine)-palladium, by the method described in by Stille, J. Amer. Chem. Soc, 1987, **109**, 5478, US Patents 4,719,218 and 5,002,941, or by using a palladium (II) catalyst in the presence of lithium chloride optionally with an added base such as triethylamine, in an inert solvent such as dimethyl formamide. Trialkyltin derivatives may be
30 conveniently obtained by metallation of the corresponding compound of Formula (IX) with a lithiating agent, such as *s*-butyl-lithium or *n*-butyllithium, in an ethereal solvent, such as tetrahydrofuran, or treatment of the bromo derivative of the corresponding compound of Formula (IX) with an alkyl lithium, followed, in each case, by treatment with a trialkyltin halide. Alternatively, the bromo- derivative of a
35 compound of Formula (IX) may be treated with a suitable heteroaryl or aryl trialkyl

tin compound in the presence of a catalyst such as *tetrakis*-(triphenyl-phosphine)-palladium, under conditions similar to those described above.


Boronic acid derivatives are also useful. Hence, a suitable derivative of a compound of Formula (IX), such as the bromo, iodo, triflate or fluorosulphonate derivative, may be reacted with a heteroaryl- or aryl-boronic acid, in the presence of
5 a palladium catalyst such as *tetrakis*-(triphenylphosphine)-palladium or $\text{PdCl}_2[\text{1,4-bis}-(\text{diphenyl-phosphino})-\text{butane}]$ in the presence of a base such as sodium bicarbonate, under reflux conditions, in a solvent such as dimethoxyethane (see Fischer and Haviniga, Rec. Trav. Chim. Pays Bas, **84**, 439, 1965, Snieckus, V.,
10 Tetrahedron Lett., **29**, 2135, 1988 and Terashimia, M., Chem. Pharm. Bull., **11**, 4755, 1985). Non-aqueous conditions, for instance, a solvent such as DMF, at a temperature of about 100°C, in the presence of a Pd(II) catalyst may also be employed (see Thompson W J *et al*, J Org Chem, **49**, 5237, 1984). Suitable boronic acid derivatives may be prepared by treating the magnesium or lithium derivative
15 with a trialkylborate ester, such as triethyl, tri-*iso*-propyl or tributylborate, according to standard procedures.

In such coupling reactions, it will be readily appreciated that due regard must be exercised with respect to functional groups present in the compounds of Formula (IX). Thus, in general, amino and sulfur substituents should be non-oxidised or
20 protected.

Compounds of Formula (IX) are imidazoles and may be obtained by any of the procedures herein before described for preparing compounds of Formula (I). In particular, an α -halo-ketone or other suitably activated ketones $\text{R}_4\text{COCH}_2\text{Hal}$ (for compounds of Formula (IX) in which T_1 is hydrogen) or $\text{R}_1\text{COCH}_2\text{Hal}$ (for
25 compounds of Formula (IX) in which T_4 is hydrogen) may be reacted with an amidine of the formula $\text{R}_2\text{NH}-\text{C}=\text{NH}$, wherein R_2 is as defined in Formula (I), or a salt thereof, in an inert solvent such as a halogenated hydrocarbon solvent, for instance chloroform, at a moderately elevated temperature, and, if necessary, in the presence of a suitable condensation agent such as a base. The preparation of suitable
30 α -halo-ketones is described in WO 91/19497. Suitable reactive esters include esters of strong organic acids such as a lower alkane sulphonic or aryl sulphonic acid, for instance, methane or *p*-toluene sulphonic acid. The amidine is preferably used as the salt, suitably the hydrochloride salt, which may then be converted into the free amidine *in situ*, by employing a two phase system in which the reactive ester is in
35 an inert organic solvent such as chloroform, and the salt is in an aqueous phase to which a solution of an aqueous base is slowly added, in dimolar amount, with

vigorous stirring. Suitable amidines may be obtained by standard methods, see for instance, Garigipati R, Tetrahedron Letters, 190, 31, 1989.

Compounds of Formula (I) may also be prepared by a process which comprises reacting a compound of Formula (IX), wherein T₁ is hydrogen, with an N-acyl heteroaryl salt, according to the method disclosed in US patent 4,803,279, US patent 4,719,218 and US patent 5,002,941, to give an intermediate in which the heteroaryl ring is attached to the imidazole nucleus and is present as a 1,4-dihydro derivative thereof, which intermediate may then be subjected to oxidative-deacylation conditions (Scheme II). The heteroaryl salt, for instance a pyridinium salt, may be either preformed or, more preferably, prepared *in situ* by adding a substituted carbonyl halide (such as an acyl halide, an aroyl halide, an arylalkyl haloformate ester, or, preferably, an alkyl haloformate ester, such as acetyl bromide, benzoylchloride, benzyl chloroformate, or, preferably, ethyl chloroformate) to a solution of the compound of Formula (IX) in the heteroaryl compound R₁H or in an inert solvent such as methylene chloride to which the heteroaryl compound has been added. Suitable deacylating and oxidising conditions are described in U.S. Patent Nos. 4,803,279, 4,719,218 and 5,002,941, which references are hereby incorporated by reference in their entirety. Suitable oxidizing systems include sulfur in an inert solvent or solvent mixture, such as decalin, decalin and diglyme, *p*-cymene, xylene or mesitylene, under reflux conditions, or, preferably, potassium *t*-butoxide in *t*-butanol with dry air or oxygen.




Scheme II

In a further process, illustrated in Scheme III below, compounds of Formula (I) may be prepared by treating a compound of Formula (X) thermally or with the aid of a cyclising agent such as phosphorus oxychloride or phosphorus pentachloride (see Engel and Steglich, Liebigs Ann Chem, 1978, 1916 and Strzybny *et al.*, J Org Chem, 1963, 28, 3381). Compounds of Formula (X) may be obtained, for instance, by acylating the corresponding α -keto-amine with an activated formate derivative such as the corresponding anhydride, under standard acylating conditions followed


by formation of the imine with R_2NH_2 . The aminoketone may be derived from the parent ketone by oximation and reduction and the requisite ketone may in turn be prepared by decarboxylation of the beta-ketoester obtained from the condensation of an aryl (heteroaryl) acetic ester with the R_1COX component.

5



Scheme III

In Scheme IV illustrated below, two (2) different routes which use ketone
 10 (formula XI) for preparing a compound of Formula (I). A heterocyclic ketone (XI) is prepared by adding the anion of the alkyl heterocycle such as 4-methyl-quinoline (prepared by treatment thereof with an alkyl lithium, such as *n*-butyl lithium) to an
 N-alkyl-O-alkoxybenzamide, ester, or any other suitably activated derivative of the
 same oxidation state. Alternatively, the anion may be condensed with a
 15 benzaldehyde, to give an alcohol which is then oxidised to the ketone (XI).



Scheme IV

20

In a further process, N-substituted compounds of Formula (I) may be prepared by treating the anion of an amide of Formula (XII):



wherein R_1 and R_2 with:

25 (a) a nitrile of the Formula (XIII):




wherein R_4 is as hereinbefore defined, or

(b) an excess of an acyl halide, for instance an acyl chloride, of the Formula (XIV):



wherein R_4 is as hereinbefore defined and Hal is halogen, or a corresponding

5 anhydride, to give a *bis*-acylated intermediate which is then treated with a source of ammonia, such as ammonium acetate.




Scheme V

10

One variation of this approach is illustrated in Scheme V above. A primary amine (R_2NH_2) is treated with a halomethyl heterocycle of Formula R_1CH_2X to give the secondary amine which is then converted to the amide by standard techniques. Alternatively the amide may be prepared as illustrated in scheme V by
 15 alkylation of the formamide with R_1CH_2X . Deprotonation of this amide with a strong amide base, such as lithium di-*iso*-propyl amide or sodium *bis*-(trimethylsilyl)amide, followed by addition of an excess of an aroyl chloride yields the *bis*-acylated compound which is then closed to an imidazole compound of Formula (I), by heating in acetic acid containing ammonium acetate. Alternatively,
 20 the anion of the amide may be reacted with a substituted aryl nitrile to produce the imidazole of Formula (I) directly.

The following description and schemes are further exemplification of the process as previously described above in Scheme I. Various pyrimidine aldehyde
 25 derivatives 6, as depicted in Scheme VI below, can be prepared by modification of the procedures of Brederick et al. (*Chem. Ber.* 1964, 97, 3407) whose disclosure is incorporated by reference herein. These pyrimidine aldehydes are then utilized as intermediates in the synthesis as further described herein.



Scheme VI

The reaction of imines with tosylmethyl isonitriles was first reported by van Leusen (van Leusen, et al., *J. Org. Chem.* **1977**, *42*, 1153.) Reported were the following conditions: tert butyl amine (*t*BuNH₂) in dimethoxyethane (DME), K₂CO₃ in MeOH, and NaH in DME. Upon re-examination of these conditions each was found to produce low yields. A second pathway involving amine exchange to produce the *t*-butyl imine followed by reaction with the isocyanide to produce a 1-*t*Bu imidazole was also operating. This will likely occur using any primary amine as a base. The secondary amines, while not preferred may be used, but may also decompose the isonitrile slowly. Reactions will likely require about 3 equivalents of amine to go to completion, resulting in approximately 50% isolated yields. Hindered secondary amines (diisopropylamine) while usable are very slow and generally not too effective. Use of tertiary and aromatic amines, such as pyridine, and triethylamine gave no reaction under certain test conditions, but more basic types such as DBU, and 4-dimethylamino pyridine (DMAP) while slow, did produce some yields and hence may be suitable for use herein.

As depicted in Schemes VII and VIII below, the pyrimidine aldehydes of Scheme VI, can be condensed with a primary amine, to generate an imine, which may suitably be isolated or reacted in situ, with the desired isonitrile in the presence of a variety of suitable bases, and solvents as described herein to afford the 5-(4-pyrimidinyl)-substituted imidazoles, wherein R₂ and R₄ are as defined herein for Formula (I) compounds.


One preferred method for preparing compounds of Formula (I) is shown below in Scheme VII, in which the imine is prepared and isolated in a separate step prior to the addition of the isonitrile. The yield for making the imines varied, and

environmentally less-acceptable solvents, such as CH_2Cl_2 were often used in their preparation.

This reaction, wherein $p=2$, requires a suitable base for the reaction to proceed. The reaction requires a base strong enough to deprotonate the isonitrile. Suitable bases include an amine, a carbonate, a hydride, or an alkyl or aryl lithium reagent; or mixtures thereof. Bases include, but are not limited to, potassium carbonate, sodium carbonate, primary and secondary amines, such as *t*-butylamine, diisopropyl amine, morpholine, piperidine, pyrrolidine, and other non-nucleophilic bases, such as DBU, DMAP and 1,4-diazabicyclo[2.2.2]octane (DABCO).

Suitable solvents for use herein, include but are not limited to the organic solvents of *N,N*-dimethyl-formamide (DMF), MeCN, halogenated solvents, such as methylene chloride or chloroform, tetrahydrofuran (THF), dimethylsulfoxide (DMSO), alcohols, such as methanol or ethanol, benzene, toluene, DME or EtOAc. Preferably the solvent is DMF, DME, THF, or MeCN, more preferably DMF.

Product isolation may generally be accomplished by adding water and filtering the product as a clean compound. In scheme VII below, Ra is as defined for Formula (I) compounds, and X is oxygen or sulfur.



Scheme VII


While not convenient for large scale work, addition of NaH to the isonitrile, perhaps with temperatures lower than 25 °C (in THF) are likely needed.

Additionally, BuLi has also been reported to be an effective base for deprotonating tosyl benzylisonitriles at -50 °C. (DiSanto, et al., *Synth. Commun.* **1995**, 25, 795).

Various temperature conditions may be utilized depending upon the preferred base. For instance, *t*BuNH₂/DME, K₂CO₃/MeOH, K₂CO₃ in DMF, at temperatures above 40 °C, the yields may drop to about 20% but little difference is expected between 0°C and 25 °C. Consequently, temperature ranges below 0°C, and above 80°C are contemplated as also being within the scope of this invention. Preferably, the temperature ranges are from about 0°C to about 25°C. For purposes


herein, room temperature, is generally depicted as 25°C, but it is recognized that this may vary from 20°C to 30°C.

- As shown in Scheme VIII below, the imine is preferably formed *in situ* in a solvent. This preferred synthesis, is a process which occurs as a one-pot synthesis.
- 5 Suitably, when the primary amine is utilized as a salt, such as in the dihydrochloride salt in the Examples, the reaction may further include a base, such as potassium carbonate, prior to the addition of the isonitrile. For hydroxy-containing amines a protected group(PG) may be required in the imine forming and cycloaddition reactions; suitably the PG is silyl (such as triethyl, diphenyl-t-butyl, dimethyl-t-butyl) or C(O)₂R, wherein R is preferably alkyl, aryl, arylalkyl moieties well known
- 10 to those skilled in the art. Reaction conditions, such as solvents, bases, temperatures, etc. are similar to those illustrated and discussed above for the isolated imine as shown in Scheme VII. One skilled in the art would readily recognize that under some circumstances, the *in situ* formation of the imine may require dehydrating
- 15 conditions, or may require acid catalysis.




Scheme VIII

- 20 Scheme IX, describes an alternative process for making compounds of formula (I). In this particular instance, the alkylthio moiety is oxidized to the methyl sulfinyl or sulfonyl moiety which is reacted with a suitable YR_a moiety.



Scheme IX

Another embodiment of the present invention is the novel hydrolysis of 2-thioalkylpyrimidine acetal to 2-thioalkylpyrimidine aldehyde, as shown in Scheme X below. Hydrolysis of the acetal to aldehyde using various known reaction conditions, such as formic acid, did not produce a satisfactory yield of the aldehyde, <13%) was obtained. The preferred synthesis involves the use of AcOH (fresh) as solvent and concentrated H₂SO₄ under heating conditions, preferably a catalytic amount of sulfuric acid. Heating conditions include temperatures from about 60 to 85°C, preferably from about 70 to about 80°C as higher temperatures show a darkening of the reaction mixture. After the reaction is completed the mixture is cooled to about room temperature and the acetic acid is removed. A more preferred alternative procedure to this involves heating the acetal in 3N HCl at 40°C for about 18 hours, cooling and extracting the bicarbonate neutralized solution into EtOAc



Scheme X

The final 2-(RaY)pyrimidin-4-yl imidazole compounds of Formula (I), as well as similar pyridine containing compounds can be prepared by one of three methods: 1) direct reaction of the 2-(RaY)pyrimidine imine with the isonitrile; 2) oxidation of the 2-alkylthiopyrimidine derivative to the corresponding sulfoxide followed by displacement with the desired HYRa under basic conditions, for example using a metal salt of HYRa or in the presence of a non-nucleophilic amine or alkali metal base; or 3) reaction of the 2-halopyrimidine or pyridine imine with the isonitrile followed by displacement with HYRa under basic conditions described in the second method, see also Adams et al., USSN 08/659,102 filed 3 June 1996, Scheme XI, whose disclosure is incorporated herein by reference in its entirety.


While these schemes herein are presented, for instance, with R₂₂ = CH₂OPh and (A) = methyl in the R₂ position, and 4-fluorophenyl for R₄, any suitable R₂ moiety or R₄ moiety may be added in this manner if it can be prepared on the primary amine. Similarly, any suitable R₄ can be added via the isonitrile route.

The compounds of Formula (II), in Scheme I, may be prepared by the methods of van Leusen et al., supra. For example, a compound of the Formula (II) may be prepared by dehydrating a compound of the Formula (IV)-Scheme I, wherein Ar, R₄ and p are as defined herein.

Suitable dehydrating agents include phosphorus oxychloride, oxalyl chloride, thionyl chloride, phosgene, or tosyl chloride in the presence of a suitable base such as triethylamine or diisopropylethylamine, or similar bases, etc. such as pyridine. Suitable solvents are dimethoxy ether, tetrahydrofuran, or halogenated solvents, preferably THF. The reaction is most efficient when the reaction temperatures are kept between -10°C and 0°C. At lower temperatures incomplete reaction occurs and at higher temperatures, the solution turns dark and the product yield drops.

The compounds of formula (IV)-Scheme I may be prepared by reacting a compound of the formula (V)-Scheme I, $R_4\text{CHO}$ where R_4 is as defined herein, with $\text{ArS(O)}_p\text{H}$ and formamide with or without water removal, preferably under dehydrating conditions, at ambient or elevated temperature e.g. 30° to 150°, conveniently at reflux, optionally in the presence of an acid catalyst. Alternatively trimethylsilylchloride can be used in place of the acid catalyst. Examples of acid catalysts include camphor-10-sulphonic acid, formic acid, p-toluenesulphonic acid, hydrogen chloride or sulphuric acid.

An optimal method of making an isonitrile of Formula (II) is illustrated below, in Scheme XI.



SCHEME XI

The conversion of the substituted aldehyde to the tosylbenzyl formamide may be accomplished by heating the aldehyde, 1-Scheme XI, with an acid, such as p-toluene-sulfonic acid, formic acid or camphorsulfonic acid; with formamide and p-toluene-sulfinic acid [under reaction conditions of about 60°C for about 24 hours].

Preferably, no solvent is used. The reaction, may give poor yields (< 30%) when solvents, such as DMF, DMSO, toluene, acetonitrile, or excess formamide are used. Temperatures less than 60°C are generally poor at producing the desired product, and temperatures in excess of 60°C may produce a product which decomposes, or
5 obtain a benzylic bis-formamide, 2-Scheme XI.

Another embodiment of the present invention is the synthesis of the tosyl benzyl formamide compound, achieved by reacting the bisformamide intermediate, 2-Scheme XI with p-toluenesulfinic acid. In this preferred route, preparation of the bis-formamide from the aldehyde is accomplished by heating the aldehyde with
10 formamide, in a suitable solvent with acid catalysis. Suitable solvents are toluene, acetonitrile, DMF, and DMSO or mixtures thereof. Acid catalysts, are those well known in the art, and include but are not limited to hydrogen chloride, p-toluenesulfonic acid, camphorsulfonic acid, and other anhydrous acids. The reaction can be conducted at temperatures ranging from about 25°C to 110°C, preferably about
15 50°C, suitably for about 4 to about 5 hours, longer reaction times are also acceptable. Product decomposition and lower yields may be observed at higher temperatures (>70°C) at prolonged reactions times. Complete conversion of the product generally requires water removal from the reaction mixture.

Preferred conditions for converting a bis-formamide derivative to the tosyl benzyl formamide are accomplished by heating the bisformamide in a suitable solvent with an acid catalyst and p-toluenesulfinic acid. Solvents for use in this reaction include but are not limited to toluene, and acetonitrile or mixtures thereof. Additional mixtures of these solvents with DMF, or DMSO may also be used but may result in lower yields. Temperatures may range from about 30°C to about
25 100°C. Temperatures lower than 40°C and higher than 60°C are not preferred as the yield and rate decreases. Preferably the range is from about 40 to 60°C, most preferably about 50°C. The optimal time is about 4 to 5 hours, although it may be longer. Preferably, acids used include but are not limited to, toluenesulfonic acid, camphorsulfonic acid, and hydrogen chloride and other anhydrous acids. Most
30 preferably the bisformamide is heated in toluene:acetonitrile in a 1:1 ratio, with p-toluenesulfinic acid and hydrogen chloride.

Another embodiment of the present invention is the preferred synthetic route for synthesis of the tosylbenzyl formamide compound which is accomplished using a one-pot procedure. This process first converts the aldehyde to the bis-formamide
35 derivative and subsequently reacts the bis-formamide derivative with toluenesulfinic acid. This procedure combines the optimized conditions into a single, efficient

process. High yields, >90% of the aryl benzylformamide may be obtained in such a manner.

Preferred reaction conditions employ a catalyst, such as trimethylsilyl chloride (TMSCl), in a preferred solvent, toluene:acetonitrile, preferably in a 1:1 ratio. A reagent, such as TMSCl, is preferred which reacts with water produced therein and at the same time produces hydrogen chloride to catalyze the reaction. Also preferred is use of hydrogen chloride and p-toluenesulfonic acid. Therefore, three suitable reaction conditions for use herein include 1) use of a dehydrating agent which also provides hydrogen chloride, such as TMSCl; or by 2) use of a suitable dehydrating agent and a suitable source of acid source, such as but not limited to, camphorsulfonic acid, hydrogen chloride or toluenesulfonic acid; and 3) alternative dehydrating conditions, such as the azeotropic removal of water, and using an acid catalyst and p-toluene sulfonic acid.

Compounds of the formula (II) where p is 2 may also be prepared by reacting in the presence of a strong base a compound of the formula (VI)-Scheme I, R_4CH_2NC with a compound of the formula (VII)-Scheme I, $ArSO_2L_1$ wherein R_4 and Ar are as defined herein and L_1 is a leaving group such as halo, e.g. fluoro. Suitable strong bases include, but are not limited to, alkyl lithiums such as butyl lithium or lithium diisopropylamide (Van Leusen *et al.*, *Tetrahedron Letters*, No. 23, 2367-68 (1972)).

The compounds of formula (VI)-Scheme I may be prepared by reacting a compound of the formula (VIII)-Scheme I, $R_4CH_2NH_2$ with an alkyl formate (e.g. ethylformate) to yield an intermediate amide which can be converted to the desired isonitrile by reacting with well known dehydrating agent, such as but not limited to oxalyl chloride, phosphorus oxychloride or tosyl chloride in the presence of a suitable base such as triethylamine.


Alternatively a compound of the formula (VIII) - Scheme I may be converted to a compound of the formula (VI)- Scheme I by reaction with chloroform and sodium hydroxide in aqueous dichloromethane under phase transfer catalysis.

The compounds of the formula (III) - Scheme I may be prepared by reacting a compound of the formula R_1CHO with a primary amine R_2NH_2 .

The amino compounds of the formula (VIII) - Scheme I are known or can be prepared from the corresponding alcohols, oximes or amides using standard functional group interconversions.

The amino compounds used to prepare the imines of formula (III) - Scheme I are known or can be prepared using standard functional group interconversions

(Scheme XII). A particularly useful and general method to prepare these amines is from the amino acids, which are readily available or if not can be prepared from the corresponding aldehyde using standard amino acid synthesis, such as the Strecker synthesis. The free amino acids or the corresponding amino protected compounds (CBZ, fMOC, or t-BOC) many of which are commercially available can be reduced to the carbinol under standard conditions. For example, borane on the carboxylic acid or if the ester, hydride agents may be employed in the reduction. The protected amino alcohols may be used as intermediates to further elaborate the side chain. Furthermore, protecting groups may be used to mask reactive functionality and thereby facilitate the formation of the imine and subsequent cycloaddition reaction to form the imidazole. An example of this is the use of a silyl protecting group on a alcohol.



Scheme XII

15

Suitable protecting groups for use with hydroxyl groups and the imidazole nitrogen are well known in the art and described in many references, for instance,

Protecting Groups in Organic Synthesis, Greene T W, Wiley-Interscience, New York, 1981. Suitable examples of hydroxyl protecting groups include silyl ethers, such as t-butyldimethyl or t-butyldiphenyl, and alkyl ethers, such as methyl connected by an alkyl chain of variable link, $(CR_{10}R_{20})_n$. Suitable examples of

5 imidazole nitrogen protecting groups include tetrahydropyranyl.

Pharmaceutically acid addition salts of compounds of Formula (I) may be obtained in known manner, for example by treatment thereof with an appropriate amount of acid in the presence of a suitable solvent.

10 METHODS OF TREATMENT

The compounds of Formula (I) or a pharmaceutically acceptable salt thereof can be used in the manufacture of a medicament for the prophylactic or therapeutic treatment of any disease state in a human, or other mammal, which is exacerbated or caused by excessive or unregulated cytokine production by such mammal's cell, such

15 as but not limited to monocytes and/or macrophages.

Compounds of Formula (I) are capable of inhibiting proinflammatory cytokines, such as IL-1, IL-6, IL-8, and TNF and are therefore of use in therapy. IL-1, IL-6, IL-8 and TNF affect a wide variety of cells and tissues and these cytokines, as well as other leukocyte-derived cytokines, are important and critical inflammatory

20 mediators of a wide variety of disease states and conditions. The inhibition of these pro-inflammatory cytokines is of benefit in controlling, reducing and alleviating many of these disease states.

Accordingly, the present invention provides a method of treating a cytokine-mediated disease which comprises administering an effective cytokine-interfering

25 amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

Compounds of Formula (I) are capable of inhibiting inducible proinflammatory proteins, such as COX-2, also referred to by many other names such as prostaglandin endoperoxide synthase-2 (PGHS-2) and are therefore of use in therapy. These proinflammatory lipid mediators of the cyclooxygenase (CO)

30 pathway are produced by the inducible COX-2 enzyme. Regulation, therefore of COX-2 which is responsible for the these products derived from arachidonic acid, such as prostaglandins affect a wide variety of cells and tissues are important and critical inflammatory mediators of a wide variety of disease states and conditions. Expression of COX-1 is not effected by compounds of Formula (I). This selective

35 inhibition of COX-2 may alleviate or spare ulcerogenic liability associated with inhibition of COX-1 thereby inhibiting prostoglandins essential for cytoprotective

effects. Thus inhibition of these pro-inflammatory mediators is of benefit in controlling, reducing and alleviating many of these disease states. Most notably these inflammatory mediators, in particular prostaglandins, have been implicated in pain, such as in the sensitization of pain receptors, or edema. This aspect of pain management therefore includes treatment of neuromuscular pain, headache, cancer pain, and arthritis pain. Compounds of Formula (I) or a pharmaceutically acceptable salt thereof, are of use in the prophylaxis or therapy in a human, or other mammal, by inhibition of the synthesis of the COX-2 enzyme.

Accordingly, the present invention provides a method of inhibiting the synthesis of COX-2 which comprises administering an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof. The present invention also provides for a method of prophylaxis treatment in a human, or other mammal, by inhibition of the synthesis of the COX-2 enzyme.

In particular, compounds of Formula (I) or a pharmaceutically acceptable salt thereof are of use in the prophylaxis or therapy of any disease state in a human, or other mammal, which is exacerbated by or caused by excessive or unregulated IL-1, IL-6, IL-8 or TNF production by such mammal's cell, such as, but not limited to, monocytes and/or macrophages.

Accordingly, in another aspect, this invention relates to a method of inhibiting the production of IL-1 in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

There are many disease states in which excessive or unregulated IL-1 production is implicated in exacerbating and/or causing the disease. These include rheumatoid arthritis, osteoarthritis, stroke, endotoxemia and/or toxic shock syndrome, other acute or chronic inflammatory disease states such as the inflammatory reaction induced by endotoxin or inflammatory bowel disease, tuberculosis, atherosclerosis, muscle degeneration, multiple sclerosis, cachexia, bone resorption, psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, gout, traumatic arthritis, rubella arthritis and acute synovitis. Recent evidence also links IL-1 activity to diabetes, pancreatic β cells disease, and Alzheimer's disease.

In a further aspect, this invention relates to a method of inhibiting the production of TNF in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

Excessive or unregulated TNF production has been implicated in mediating or exacerbating a number of diseases including rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, adult
5 respiratory distress syndrome, stroke, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcoisosis, bone resorption diseases, such as osteoporosis, reperfusion injury, graft vs. host reaction, allograft rejections, fever and myalgias due to infection, such as influenza, cachexia secondary to infection or malignancy, cachexia secondary to acquired immune deficiency
10 syndrome (AIDS), AIDS, ARC (AIDS related complex), keloid formation, scar tissue formation, Crohn's disease, ulcerative colitis and pyresis.

Compounds of Formula (I) are also useful in the treatment of viral infections, where such viruses are sensitive to upregulation by TNF or will elicit TNF production *in vivo*. The viruses contemplated for treatment herein are those that
15 produce TNF as a result of infection, or those which are sensitive to inhibition, such as by decreased replication, directly or indirectly, by the TNF inhibiting-compounds of Formula (I). Such viruses include, but are not limited to HIV-1, HIV-2 and HIV-3, Cytomegalovirus (CMV), Influenza, adenovirus and the Herpes group of viruses, such as but not limited to, Herpes Zoster and Herpes Simplex. Accordingly, in a
20 further aspect, this invention relates to a method of treating a mammal afflicted with a human immunodeficiency virus (HIV) which comprises administering to such mammal an effective TNF inhibiting amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

Compounds of Formula (I) may also be used in association with the
25 veterinary treatment of mammals, other than in humans, in need of inhibition of TNF production. TNF mediated diseases for treatment, therapeutically or prophylactically, in animals include disease states such as those noted above, but in particular viral infections. Examples of such viruses include, but are not limited to, lentivirus infections such as, equine infectious anaemia virus, caprine arthritis virus, visna virus, or maedi virus or retrovirus infections, such as but not limited to feline
30 immunodeficiency virus (FIV), bovine immunodeficiency virus, or canine immunodeficiency virus or other retroviral infections.

The compounds of Formula (I) may also be used topically in the treatment or prophylaxis of topical disease states mediated by or exacerbated by excessive
35 cytokine production, such as by IL-1 or TNF respectively, such as inflamed joints, eczema, psoriasis and other inflammatory skin conditions such as sunburn;

inflammatory eye conditions including conjunctivitis; pyresis, pain and other conditions associated with inflammation.

Compounds of Formula (I) have also been shown to inhibit the production of IL-8 (Interleukin-8, NAP). Accordingly, in a further aspect, this invention relates to a method of inhibiting the production of IL-8 in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

There are many disease states in which excessive or unregulated IL-8 production is implicated in exacerbating and/or causing the disease. These diseases are characterized by massive neutrophil infiltration such as, psoriasis, inflammatory bowel disease, asthma, cardiac and renal reperfusion injury, adult respiratory distress syndrome, thrombosis and glomerulonephritis. All of these diseases are associated with increased IL-8 production which is responsible for the chemotaxis of neutrophils into the inflammatory site. In contrast to other inflammatory cytokines (IL-1, TNF, and IL-6), IL-8 has the unique property of promoting neutrophil chemotaxis and activation. Therefore, the inhibition of IL-8 production would lead to a direct reduction in the neutrophil infiltration.

The compounds of Formula (I) are administered in an amount sufficient to inhibit cytokine, in particular IL-1, IL-6, IL-8 or TNF, production such that it is regulated down to normal levels, or in some case to subnormal levels, so as to ameliorate or prevent the disease state. Abnormal levels of IL-1, IL-6, IL-8 or TNF, for instance in the context of the present invention, constitute: (i) levels of free (not cell bound) IL-1, IL-6, IL-8 or TNF greater than or equal to 1 picogram per ml; (ii) any cell associated IL-1, IL-6, IL-8 or TNF; or (iii) the presence of IL-1, IL-6, IL-8 or TNF mRNA above basal levels in cells or tissues in which IL-1, IL-6, IL-8 or TNF, respectively, is produced.

The discovery that the compounds of Formula (I) are inhibitors of cytokines, specifically IL-1, IL-6, IL-8 and TNF is based upon the effects of the compounds of Formulas (I) on the production of the IL-1, IL-8 and TNF in *in vitro* assays which are described herein.

As used herein, the term "inhibiting the production of IL-1 (IL-6, IL-8 or TNF)" refers to:

a) a decrease of excessive *in vivo* levels of the cytokine (IL-1, IL-6, IL-8 or TNF) in a human to normal or sub-normal levels by inhibition of the *in vivo* release of the cytokine by all cells, including but not limited to monocytes or macrophages;

b) a down regulation, at the genomic level, of excessive *in vivo* levels of the cytokine (IL-1, IL-6, IL-8 or TNF) in a human to normal or sub-normal levels;

c) a down regulation, by inhibition of the direct synthesis of the cytokine (IL-1, IL-6, IL-8 or TNF) as a postranslational event; or

5 d) a down regulation, at the translational level, of excessive *in vivo* levels of the cytokine (IL-1, IL-6, IL-8 or TNF) in a human to normal or sub-normal levels.

As used herein, the term "TNF mediated disease or disease state" refers to any and all disease states in which TNF plays a role, either by production of TNF itself, or by TNF causing another monokine to be released, such as but not limited to
10 IL-1, IL-6 or IL-8. A disease state in which, for instance, IL-1 is a major component, and whose production or action, is exacerbated or secreted in response to TNF, would therefore be considered a disease state mediated by TNF.

As used herein, the term "cytokine" refers to any secreted polypeptide that affects the functions of cells and is a molecule which modulates interactions between
15 cells in the immune, inflammatory or hematopoietic response. A cytokine includes, but is not limited to, monokines and lymphokines, regardless of which cells produce them. For instance, a monokine is generally referred to as being produced and secreted by a mononuclear cell, such as a macrophage and/or monocyte. Many other cells however also produce monokines, such as natural killer cells, fibroblasts,
20 basophils, neutrophils, endothelial cells, brain astrocytes, bone marrow stromal cells, epidermal keratinocytes and B-lymphocytes. Lymphokines are generally referred to as being produced by lymphocyte cells. Examples of cytokines include, but are not limited to, Interleukin-1 (IL-1), Interleukin-6 (IL-6), Interleukin-8 (IL-8), Tumor
Necrosis Factor-alpha (TNF-a) and Tumor Necrosis Factor beta (TNF-β).

25 As used herein, the term "cytokine interfering" or "cytokine suppressive amount" refers to an effective amount of a compound of Formula (I) which will cause a decrease in the *in vivo* levels of the cytokine to normal or sub-normal levels, when given to a patient for the prophylaxis or treatment of a disease state which is exacerbated by, or caused by, excessive or unregulated cytokine production.

30 As used herein, the cytokine referred to in the phrase "inhibition of a cytokine, for use in the treatment of a HIV-infected human" is a cytokine which is implicated in (a) the initiation and/or maintenance of T cell activation and/or activated T cell-mediated HIV gene expression and/or replication and/or (b) any cytokine-mediated disease associated problem such as cachexia or muscle degeneration.

35 As TNF-β (also known as lymphotoxin) has close structural homology with TNF-a (also known as cachectin) and since each induces similar biologic responses

and binds to the same cellular receptor, both TNF- α and TNF- β are inhibited by the compounds of the present invention and thus are herein referred to collectively as "TNF" unless specifically delineated otherwise.

A new member of the MAP kinase family, alternatively termed CSBP, p38, or RK, has been identified independently by several laboratories. Activation of this novel protein kinase via dual phosphorylation has been observed in different cell systems upon stimulation by a wide spectrum of stimuli, such as physicochemical stress and treatment with lipopolysaccharide or proinflammatory cytokines such as interleukin-1 and tumor necrosis factor. The cytokine biosynthesis inhibitors, of the present invention, compounds of Formula (I) have been determined to be potent and selective inhibitors of CSBP/p38/RK kinase activity. These inhibitors are of aid in determining the signaling pathways involvement in inflammatory responses. In particular, for the first time a definitive signal transduction pathway can be prescribed to the action of lipopolysaccharide in cytokine production in macrophages. In addition to those diseases already noted, treatment of stroke, neurotrauma, cardiac and renal reperfusion injury, congestive heart failure, chronic renal failure, angiogenesis & related processes, such as cancer, thrombosis, glomerulonephritis, diabetes and pancreatic b cells, multiple sclerosis, muscle degeneration, eczema, psoriasis, sunburn, and conjunctivitis are also included.

The CSBP inhibitors were subsequently tested in a number of animal models for anti-inflammatory activity. Model systems were chosen that were relatively insensitive to cyclooxygenase inhibitors in order to reveal the unique activities of cytokine suppressive agents. The inhibitors exhibited significant activity in many such in vivo studies. Most notable are its effectiveness in the collagen-induced arthritis model and inhibition of TNF production in the endotoxic shock model. In the latter study, the reduction in plasma level of TNF correlated with survival and protection from endotoxic shock related mortality. Also of great importance are the compounds effectiveness in inhibiting bone resorption in a rat fetal long bone organ culture system. Griswold et al., (1988) *Arthritis Rheum.* **31**:1406-1412; Badger, et al., (1989) *Circ. Shock* **27**, 51-61; Votta et al., (1994) *in vitro. Bone* **15**, 533-538; Lee et al., (1993). *B Ann. N. Y. Acad. Sci.* **696**, 149-170.

Chronic diseases which have an inappropriate angiogenic component are various ocular neovascularizations, such as diabetic retinopathy and macular degeneration. Other chronic diseases which have an excessive or increased proliferation of vasculature are tumor growth and metastasis, atherosclerosis, and

certain arthritic conditions. Therefore CSBP kinase inhibitors will be of utility in the blocking of the angiogenic component of these disease states.

The term "excessive or increased proliferation of vasculature inappropriate angiogenesis" as used herein includes, but is not limited to, diseases which are
5 characterized by hemangiomas and ocular diseases.

The term "inappropriate angiogenesis" as used herein includes, but is not limited to, diseases which are characterized by vesicle proliferation with accompanying tissue proliferation, such as occurs in cancer, metastasis, arthritis and atherosclerosis.

10 Accordingly, the present invention provides a method of treating a CSBP kinase mediated disease in a mammal in need thereof, preferably a human, which comprises administering to said mammal, an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

It has now been found that the branching of the R₂ moiety, such as in the R₂₂
15 term provides for improved activity against the CSBP enzyme, and for improved *in vivo* activity versus the unbranched R₂ alkyl chain. such as disclosed in US Patent 5,593,992.

In order to use a compound of Formula (I) or a pharmaceutically acceptable salt thereof in therapy, it will normally be formulated into a pharmaceutical
20 composition in accordance with standard pharmaceutical practice. This invention, therefore, also relates to a pharmaceutical composition comprising an effective, non-toxic amount of a compound of Formula (I) and a pharmaceutically acceptable carrier or diluent.

Compounds of Formula (I), pharmaceutically acceptable salts thereof and
25 pharmaceutical compositions incorporating such may conveniently be administered by any of the routes conventionally used for drug administration, for instance, orally, topically, parenterally or by inhalation. The compounds of Formula (I) may be administered in conventional dosage forms prepared by combining a compound of Formula (I) with standard pharmaceutical carriers according to conventional
30 procedures. The compounds of Formula (I) may also be administered in conventional dosages in combination with a known, second therapeutically active compound. These procedures may involve mixing, granulating and compressing or dissolving the ingredients as appropriate to the desired preparation. It will be appreciated that the form and character of the pharmaceutically acceptable character
35 or diluent is dictated by the amount of active ingredient with which it is to be combined, the route of administration and other well-known variables. The carrier(s)

must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

The pharmaceutical carrier employed may be, for example, either a solid or liquid. Exemplary of solid carriers are lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid and the like. Exemplary of liquid
5 carriers are syrup, peanut oil, olive oil, water and the like. Similarly, the carrier or diluent may include time delay material well known to the art, such as glyceryl mono-stearate or glyceryl distearate alone or with a wax.

A wide variety of pharmaceutical forms can be employed. Thus, if a solid
10 carrier is used, the preparation can be tableted, placed in a hard gelatin capsule in powder or pellet form or in the form of a troche or lozenge. The amount of solid carrier will vary widely but preferably will be from about 25mg. to about 1g. When a liquid carrier is used, the preparation will be in the form of a syrup, emulsion, soft gelatin capsule, sterile injectable liquid such as an ampule or nonaqueous liquid
15 suspension.

Compounds of Formula (I) may be administered topically, that is by non-systemic administration. This includes the application of a compound of Formula (I) externally to the epidermis or the buccal cavity and the instillation of such a compound into the ear, eye and nose, such that the compound does not significantly
20 enter the blood stream. In contrast, systemic administration refers to oral, intravenous, intraperitoneal and intramuscular administration.

Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin to the site of inflammation such as liniments, lotions, creams, ointments or pastes, and drops suitable for
25 administration to the eye, ear or nose. The active ingredient may comprise, for topical administration, from 0.001% to 10% w/w, for instance from 1% to 2% by weight of the formulation. It may however comprise as much as 10% w/w but preferably will comprise less than 5% w/w, more preferably from 0.1% to 1% w/w of the formulation.

30 Lotions according to the present invention include those suitable for application to the skin or eye. An eye lotion may comprise a sterile aqueous solution optionally containing a bactericide and may be prepared by methods similar to those for the preparation of drops. Lotions or liniments for application to the skin may also include an agent to hasten drying and to cool the skin, such as an alcohol or
35 acetone, and/or a moisturizer such as glycerol or an oil such as castor oil or arachis oil.

Creams, ointments or pastes according to the present invention are semi-solid Formulations of the active ingredient for external application. They may be made by mixing the active ingredient in finely-divided or powdered form, alone or in solution or suspension in an aqueous or non-aqueous fluid, with the aid of suitable

5 machinery, with a greasy or non-greasy base. The base may comprise hydrocarbons such as hard, soft or liquid paraffin, glycerol, beeswax, a metallic soap; a mucilage; an oil of natural origin such as almond, corn, arachis, castor or olive oil; wool fat or its derivatives or a fatty acid such as steric or oleic acid together with an alcohol such as propylene glycol or a macrogel. The formulation may incorporate any suitable
10 surface active agent such as an anionic, cationic or non-ionic surfactant such as a sorbitan ester or a polyoxyethylene derivative thereof. Suspending agents such as natural gums, cellulose derivatives or inorganic materials such as siliceous silicas, and other ingredients such as lanolin, may also be included.

Drops according to the present invention may comprise sterile aqueous or
15 oily solutions or suspensions and may be prepared by dissolving the active ingredient in a suitable aqueous solution of a bactericidal and/or fungicidal agent and/or any other suitable preservative, and preferably including a surface active agent. The resulting solution may then be clarified by filtration, transferred to a suitable container which is then sealed and sterilized by autoclaving or maintaining
20 at 98-100°C. for half an hour. Alternatively, the solution may be sterilized by filtration and transferred to the container by an aseptic technique. Examples of bactericidal and fungicidal agents suitable for inclusion in the drops are phenylmercuric nitrate or acetate (0.002%), benzalkonium chloride (0.01%) and chlorhexidine acetate (0.01%). Suitable solvents for the preparation of an oily
25 solution include glycerol, diluted alcohol and propylene glycol.

Compounds of Formula (I) may be administered parenterally, that is by intravenous, intramuscular, subcutaneous intranasal, intrarectal, intravaginal or intraperitoneal administration. The subcutaneous and intramuscular forms of parenteral administration are generally preferred. Appropriate dosage forms for such
30 administration may be prepared by conventional techniques. Compounds of Formula (I) may also be administered by inhalation, that is by intranasal and oral inhalation administration. Appropriate dosage forms for such administration, such as an aerosol formulation or a metered dose inhaler, may be prepared by conventional techniques.

35 For all methods of use disclosed herein for the compounds of Formula (I), the daily oral dosage regimen will preferably be from about 0.1 to about 80 mg/kg of

total body weight, preferably from about 0.2 to 30 mg/kg, more preferably from about 0.5 mg to 15mg. The daily parenteral dosage regimen about 0.1 to about 80 mg/kg of total body weight, preferably from about 0.2 to about 30 mg/kg, and more preferably from about 0.5 mg to 15mg/kg. The daily topical dosage regimen will preferably be from 0.1 mg to 150 mg, administered one to four, preferably two or three times daily. The daily inhalation dosage regimen will preferably be from about 0.01 mg/kg to about 1 mg/kg per day. It will also be recognized by one of skill in the art that the optimal quantity and spacing of individual dosages of a compound of Formula (I) or a pharmaceutically acceptable salt thereof will be determined by the nature and extent of the condition being treated, the form, route and site of administration, and the particular patient being treated, and that such optimums can be determined by conventional techniques. It will also be appreciated by one of skill in the art that the optimal course of treatment, i.e., the number of doses of a compound of Formula (I) or a pharmaceutically acceptable salt thereof given per day for a defined number of days, can be ascertained by those skilled in the art using conventional course of treatment determination tests.

The novel compounds of Formula (I) may also be used in association with the veterinary treatment of mammals, other than humans, in need of inhibition of CSBP/p38 or cytokine inhibition or production. In particular, CSBP/p38 mediated diseases for treatment, therapeutically or prophylactically, in animals include disease states such as those noted herein in the Methods of Treatment section, but in particular viral infections. Examples of such viruses include, but are not limited to, lentivirus infections such as, equine infectious anaemia virus, caprine arthritis virus, visna virus, or maedi virus or retrovirus infections, such as but not limited to feline immunodeficiency virus (FIV), bovine immunodeficiency virus, or canine immunodeficiency virus or other retroviral infections.

The invention will now be described by reference to the following biological examples which are merely illustrative and are not to be construed as a limitation of the scope of the present invention.

BIOLOGICAL EXAMPLES

The cytokine-inhibiting effects of compounds of the present invention may be determined by the following *in vitro* assays:

Assays for Interleukin - 1 (IL-1), Interleukin -8 (IL-8), and Tumour Necrosis Factor (TNF) are well known in the art, and may be found in a number of publications,

and patents. Representative suitable assays for use herein are described in Adams et al., US 5,593,992, whose disclosure is incorporated by reference in its entirety.

In vivo TNF assay:

5

(1) Griswold *et al.*, Drugs Under Exp. and Clinical Res., XIX (6), 243-248 (1993); or

(2) Boehm, *et al.*, *Journal Of Medicinal Chemistry* 39, 3929-3937 (1996) whose disclosures are incorporated by reference herein in their entirety.

10

LPS-induced TNF α Production in Mice and Rats

In order to evaluate in vivo inhibition of LPS-induced TNF α production in rodents, both mice and rats are injected with LPS.

Mouse Method

15

Male Balb/c mice from Charles River Laboratories are pretreated (30 minutes) with compound or vehicle. After the 30 min. pretreat time, the mice are given LPS (lipopolysaccharide from *Esheria coli* Serotype 055-85, Sigma Chemical Co., St Louis, MO) 25 ug/mouse in 25 ul phosphate buffered saline (pH 7.0) intraperitoneally. Two hours later the mice are killed by CO₂ inhalation and blood samples are collected by exsanguination into heparinized blood collection tubes and stored on ice. The blood samples are centrifuged and the plasma collected and stored at -20°C until assayed for TNF α by ELISA.

20

Rat Method

25

Male Lewis rats from Charles River Laboratories are pretreated at various times with compound or vehicle. After a determined pretreat time, the rats are given LPS (lipopolysaccharide from *Esheria coli* Serotype 055-85, Sigma Chemical Co., St Louis, MO) 3.0 mg/kg intraperitoneally. The rats are killed by CO₂ inhalation and heparinized whole blood is collected from each rat by cardiac puncture 90 minutes after the LPS injection. The blood samples are centrifuged and the plasma collected for analysis by ELISA for TNF α levels.

30

ELISA Method

TNF α levels were measured using a sandwich ELISA, as described in Olivera et al., *Circ. Shock*, 37, 301-306, (1992), whose disclosure is incorporated by reference in its entirety herein, using a hamster monoclonal antimurine TNF α (Genzyme, Boston, MA) as the capture antibody and a polyclonal rabbit antimurine TNF α

35

(Genzyme) as the second antibody. For detection, a peroxidase-conjugated goat antirabbit antibody (Pierce, Rockford, IL) was added, followed by a substrate for peroxidase (1 mg/ml orthophenylenediamine with 1% urea peroxide). TNF α levels in the plasma samples from each animal were calculated from a standard curve generated with recombinant murine TNF α (Genzyme).

LPS-Stimulated Cytokine Production in Human Whole Blood

Assay: Test compound concentrations were prepared at 10 X concentrations and LPS prepared at 1 ug/ml (final conc. of 50 ng/ml LPS) and added in 50 uL volumes to 1.5 mL eppendorf tubes. Heparinized human whole blood was obtained from healthy volunteers and was dispensed into eppendorf tubes containing compounds and LPS in 0.4 mL volumes and the tubes incubated at 37 C. Following a 4 hour incubation, the tubes were centrifuged at 5000 rpm for 5 minutes in a TOMY microfuge, plasma was withdrawn and frozen at -80 C.

Cytokine measurement: IL-1 and/or TNF were quantified using a standardized ELISA technology. An in-house ELISA kit was used to detect human IL-1 and TNF. Concentrations of IL-1 or TNF were determined from standard curves of the appropriate cytokine and IC50 values for test compound (concentration that inhibited 50% of LPS-stimulated cytokine production) were calculated by linear regression analysis.

CSBP/p38 Kinase Assay:

This assay measures the CSBP/p38-catalyzed transfer of ^{32}P from [α - ^{32}P]ATP to threonine residue in an epidermal growth factor receptor (EGFR)-derived peptide (T669) with the following sequence: KRELVEPLTPSGEAPNQALLR (residues 661-681). (See Gallagher *et al.*, "Regulation of Stress Induced Cytokine Production by Pyridinyl Imidazoles: Inhibition of CSBP Kinase", BioOrganic & Medicinal Chemistry, 1997, 5, 49-64).

Reactions were carried in round bottom 96 well plate (from Corning) in a 30 ml volume. Reactions contained (in final concentration): 25 mM Hepes, pH7.5; 8 mM MgCl₂; 0.17 mM ATP (the K_m[ATP] of p38 (see Lee *et al.*, Nature 300, n72 pg 639-746 (Dec. 1994)); 2.5 uCi of [γ - ^{32}P]ATP; 0.2 mM sodium orthovanadate; 1 mM DTT; 0.1% BSA; 10% glycerol; 0.67 mM T669 peptide; and 2-4 nM of yeast-expressed, activated and purified p38. Reactions were initiated by the addition of [γ - ^{32}P]Mg/ATP, and incubated for 25 min. at 37 °C. Inhibitors (dissolved in DMSO) were incubated with the reaction mixture on ice for 30 minutes prior to

adding the ^{32}P -ATP. Final DMSO concentration was 0.16%. Reactions were terminated by adding 10 μl of 0.3 M phosphoric acid, and phosphorylated peptide was isolated from the reactions by capturing it on p81 phosphocellulose filters. Filters were washed with 75 mM phosphoric acids, and incorporated ^{32}P was

5 quantified using beta scintillation counter. Under these conditions, the specific activity of p38 was 400-450 pmol/pmol enzyme, and the activity was linear for up to 2 hr of incubation. The kinase activity values were obtained after subtracting values generated in the absence of substrate which were 10-15% of total values.

Representative final compounds of Formula (I), Examples 1 to 23 have

10 demonstrated positive inhibitory activity of an IC_{50} of $< 50\mu\text{M}$ in this binding assay or a similar assay.

Prostaglandin endoperoxide synthase-2 (PGHS-2) assay:

This assay describes a method for determining the inhibitory effects of

15 compounds of Formula (I) on human PGHS-2 protein expression in LPS stimulated human monocytes. A suitable assay for PGHS-2 protein expression may be found in a number of publications, including US Patent 5,593,992 whose disclosure is incorporated herein by reference.

20 **TNF- α in Traumatic Brain Injury Assay**

This assay provides for examination of the expression of tumor necrosis factor mRNA in specific brain regions which follow experimentally induced lateral fluid-percussion traumatic brain injury (TBI) in rats. Since TNF- α is able to induce nerve growth factor (NGF) and stimulate the release of other cytokines from activated

25 astrocytes, this post-traumatic alteration in gene expression of TNF- α plays an important role in both the acute and regenerative response to CNS trauma. A suitable assay may be found in WO 97/35856 whose disclosure is incorporated herein by reference.

30 **CNS Injury model for IL- β mRNA**

This assay characterizes the regional expression of interleukin- 1β (IL- 1β) mRNA in specific brain regions following experimental lateral fluid-percussion traumatic brain injury (TBI) in rats. Results from these assays indicate that following TBI, the temporal expression of IL- 1β mRNA is regionally stimulated in specific brain

35 regions. These regional changes in cytokines, such as IL- 1β play a role in the post-

traumatic pathologic or regenerative sequelae of brain injury. A suitable assay may be found in WO 97/35856 whose disclosure is incorporated herein by reference.

Angiogenesis Assay:

- 5 Described in WO 97/32583, whose disclosure is incorporated herein by reference, is an assay for determination of inflammatory angiogenesis which may be used to show that cytokine inhibition will stop the tissue destruction of excessive or inappropriate proliferation of blood vessels.

10 SYNTHETIC EXAMPLES

- The invention will now be described by reference to the following examples which are merely illustrative and are not to be construed as a limitation of the scope of the present invention. All temperatures are given in degrees centigrade, all solvents are highest available purity and all reactions run under anhydrous conditions
15 in an argon atmosphere unless otherwise indicated.

- In the Examples, all temperatures are in degrees Centigrade (°C). Mass spectra were performed upon a VG Zab mass spectrometer using fast atom bombardment or on a micromass platform electrospray ionization mass spectrometer in the positive ion mode using 95:5 CH₃CN/CH₃OH with 1% formic acid as the
20 carrier solvent, unless otherwise indicated. ¹H-NMR (hereinafter "NMR") spectra were recorded at 250 MHz using a Bruker AM 250 or Am 400 spectrometer. Multiplicities indicated are: s=singlet, d=doublet, t=triplet, q=quartet, m=multiplet and br indicates a broad signal. Sat. indicates a saturated solution, eq indicates the proportion of a molar equivalent of reagent relative to the principal reactant.

- 25 Flash chromatography is run over Merck Silica gel 60 (230 - 400 mesh).

Example 1

(S)-1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-fluorophenoxy)pyrimidin-4-yl]imidazole

- 30 a) 4-Fluorophenyl-tolylsulfonmethylformamide

- To a suspension of p-toluenesulfonic acid sodium salt (30 grams, hereinafter "g") in H₂O (100 milliliters (hereinafter "mL")) was added methyl t-butyl ether (50 mL) followed by dropwise addition of conc. HCl (15 mL). After stirring 5 min., the organic phase was removed and the aqueous phase was extracted with methyl t-butyl ether. The organic phase was dried (Na₂SO₄) and concentrated to near
35 dryness. Hexane was added and the free acid was filtered. The p-toluenesulfonic

acid (22 g, 140.6 millimole (hereinafter "mmol")), p-fluorobenzaldehyde (22 mL, 206 mmol), formamide (20 mL, 503 mmol) and camphor sulphonic acid (4 g, 17.3 mmol) were combined and stirred at 60°C for about 18 hours (hereinafter "h").

The resulting solid was broken up and stirred with a mixture of MeOH (35 mL) and
5 hexane (82 mL) then filtered. The solid was resuspended in MeOH/hexane (1:3, 200 mL) and stirred vigorously to break up remaining chunks. Filtration afforded the title compound (27 g, 62 % yield). ¹H NMR (400 MHz, CDCl₃): d 8.13 (s, 1H), 7.71 (d, 2H), 7.43 (dd, 2H), 7.32 (d, 2H), 7.08 (t, 2H), 6.34 (d, 1H), 2.45 (s, 3H).

10 b) 4-Fluorophenyl-tolylsulfonomethylisocyanide

The compound in the previous step (2.01g, 6.25 mmol) in ethyleneglycol dimethylether (DME) (32 mL) was cooled to -10°C. POCl₃ (1.52 mL, 16.3 mmol) was added followed by the dropwise addition of triethylamine (4.6 mL, 32.6 mmol) in DME (3mL) keeping the internal temperature below -5°C. The mixture was
15 gradually warmed over 1 h., quenched in H₂O and extracted with EtOAc. The organic phase was washed with saturated aqueous NaHCO₃, dried (Na₂SO₄), and concentrated. The resulting residue was triturated with petroleum ether and filtered to afford the title compound (1.7 g, 90% yield). ¹H NMR (CDCl₃): d 7.63 (d, 2H), 7.33 (m, 4H), 7.10 (t, 2H), 5.60 (s, 1H), 2.50 (s, 3H)

20 c) 2-Propylthiopyrimidine-4-carboxaldehyde dimethyl acetal

Charge a 1 L 3-necked flask equipped with a stir bar, thermometer, 100 mL addition funnel and reflux condensor with N,N-dimethylformamide dimethyl acetal (88.7 g, 98.9 mL, 700 mmol) and pyruvaldehyde dimethyl acetal (85.3 g, 86.8 mL, 700 mmol) and heat in an oil bath at 110 °C for 3-4 h. Cool the solution to 85 °C
25 and add thiourea (48.9 g, 636.4 mmol) and NaOMe (25 wt % in MeOH, 151.2 g, 160 mL, 700 mmol) and stir at 85 °C for 3-4 h. Cool the solution to 65 °C and charge 1-bromopropane (86.9 g, 64.4 mL, 700 mmol) to the addition funnel and add slowly over 10-15 min to the reaction, bringing the solution to a mild reflux. After 1 h, add 100 mL of EtOAc to the reaction and bring the oil bath temperature to 95 °C.
30 Replace the reflux condensor with a distillation head and distill 150-200 mL of solvent from the reaction. Add an additional 400 mL of EtOAc and 120 mL of H₂O and stir at 50 °C for 5 min. Transfer to a separatory funnel and separate the aqueous phase. Add 60 mL of H₂O, agitate, and separate the aqueous phase. A sample was concentrated to give a yellow oil: ¹H NMR (300 MHz, CDCl₃) d 8.53 (1H, d, J 5.0 Hz), 7.16 (1H, d, J = 5.0 Hz), 5.17 (1H, s), 3.42 (3H, s), 3.14 (2H, t, J = 7.3 Hz),
35 1.76 (2H, m), 1.05 (3H, t, J = 7.3 Hz).

Alternatively, bromopropane can be replaced with any suitable alkyl halide and the alkylation process can occur at about 0 to about 100 °C.

d) 2-Propylthiopyrimidine-4-carboxaldehyde

The product of the previous step (24 g, 105 mmol) was dissolved in THF (75 mL) and 3N HCL (150 mL) was added. The resulting mixture was stirred under argon and heated to 57 °C for 4 h. The THF was stripped off and the mixture was cooled in an ice bath. EtOAc (300 mL) was added followed by the addition of solid NaHCO₃. Additional H₂O was added to dissolve all the solid, and the aqueous phase was extracted with EtOAc (3 x 150 mL). The organic phases were combined, dried (Na₂SO₄) and concentrated to give a brown oil. The crude product was purified by flash chromatography (silica gel, 0-1% MeOH/CH₂Cl₂) to give the title compound as a yellow oil. ¹H NMR (400 MHz, CDCl₃): δ 9.95 (s, 1H), 8.78 (d, 1H), 7.45 (d, 1H), 3.21 (t, 2H), 1.82 (m, 2H), 1.1 (t, 3H).

e) 2-Propylthiopyrimidine-4-carboxaldehyde[(S)-2-amino-1-propanol]imine

To a solution of 2-propylthiopyrimidine-4-carboxaldehyde (10.9 g, 60 mmol) in CH₂Cl₂ (200 mL) was added (S)-2 amino-1-propanol (5.85 g, 78 mmol). The solution was stirred at room temperature under argon for 16 h. The solution was concentrated to give the title compound. ES (+) MS m/e = 240 (MH⁺)

f) (S)-1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(propylthio)pyrimidin-4-yl]imidazole

The product of the previous step (14.7 g, ~60 mmol) was dissolved in DMF (200 mL) and stirred under argon. Potassium carbonate (6.6 g, 48 mmol) was added followed by the addition of the product of example 1(b) (12.14 g, 42 mmol). The mixture was stirred at rt for 72 h. The DMF was pumped off and the residue was partitioned between EtOAc and water. The organic phase was separated, washed with brine, dried (Na₂SO₄), and concentrated. The crude product was purified by flash chromatography (silica gel, 0-4% MeOH/CH₂Cl₂) to give the title compound as a yellow solid. ES (+) MS m/e = 373 (MH⁺)

g) (S)-1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(propylsulfonyl)-pyrimidin-4-yl]imidazole

The product of example 1(f) (4g, 10.75mmol) was dissolved in methanol (100 mL) and cooled in an ice bath while stirring under argon. OXONE (8.26g, 13.44) mmol in H₂O (60 mL) was added and the mixture was stirred at rt for 12 h. The MeOH was stripped and the residue partitioned between EtOAc and H₂O. The mixture was made basic by the addition of solid K₂CO₃ and extracted with EtOAc.

the combined organic extracts were washed with brine, dried (Na_2SO_4), and concentrated to give the title compound as a yellow solid.

ES (+) MS $m/e = 405$ (MH^+)

h) (S)-1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-fluorophenoxy)-pyrimidin-4-yl]imidazole

NaH (95%) (252 mg, 10 mmol) was added in small portions to a solution of 4-fluorophenol (2.21 g, 19.8 mmol) in dry THF (50 mL). After the vigorous reaction had subsided, this solution was added to a solution of the product of example 1(g) (2 g, 4.95 mmol) dissolved in dry THF (200 mL). The reaction mixture was stirred at rt under argon for 1 h. The reaction mixture was concentrated, and the residue partitioned between EtOAc and H_2O . The organic phase was washed with 1N NaOH, brine, dried (Na_2SO_4), and concentrated. The crude product was purified by flash chromatography (silica gel, 0-4% MeOH/ CH_2Cl_2) to give the title compound as a light yellow solid.

ES (+) MS $m/e = 409$ (MH^+)

Example 2

(R)-1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-fluorophenoxy)pyrimidin-4-yl]imidazole

Following the procedures of examples 1 (e) - (h) except using (R)-2 amino-1-propanol in place of (S)-2 amino-1-propanol in step 1 (e) afforded the title compound as an off-white solid. ES (+) MS $m/e = 409$ (MH^+)

Example 3

1-(1,3-Dihydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-fluorophenoxy)pyrimidin-4-yl]imidazole

Following the procedures of examples 1 (e) - (h) except using 2-amino-1,3-propanediol in place of (S)-2 amino-1-propanol in step 1 (e) afforded the title compound as an off-white solid. ES (+) MS $m/e = 425$ (MH^+)

Example 4

(R)-1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy)pyrimidin-4-yl]imidazole

Following the procedures of examples 1 (e) - (h) except using (R)-2 amino-1-propanol in place of (S)-2 amino-1-propanol in step 1 (e) and using phenol in place

of 4-fluorophenol in step 1 (h) afforded the title compound as light yellow crystals.
ES (+) MS m/e = 391 (MH⁺)

Example 5

5 (S)-1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy-
pyrimidin-4-yl)imidazole

Following the procedure of example 1 (h) except using phenol in place of 4-fluorophenol, afforded the title compound as light yellow crystals.

ES (+) MS m/e = 391 (MH⁺)

Example 6

10

(+/-)-1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy-
pyrimidin-4-yl)imidazole

Following the procedures of examples 1 (e) - (h) except using (+/-)-2 amino-1-propanol in place of (S)-2 amino-1-propanol in step 1 (e) and using phenol in place
15 of 4-fluorophenol in step 1 (h) afforded the title compound as light yellow crystals.

ES (+) MS m/e = 391 (MH⁺)

Example 7

20

(+/-)-1-(1-Phenoxyprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy-
pyrimidin-4-yl)imidazole

Following the procedures of examples 1 (e) - (h) except using (+/-)-1-phenoxy-2-aminopropane in place of (S)-2 amino-1-propanol in step 1 (e) and using phenol in place of 4-fluorophenol in step 1 (h) afforded the title compound as a light yellow solid. ES (+) MS m/e = 467 (MH⁺)

25

Example 8

(R)-1-(1-Hydroxy-3-phenylprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy-
pyrimidin-4-yl)imidazole

Following the procedures of examples 1 (e) - (h) except using (R)-2 amino-3-phenyl-1-propanol in place of (S)-2 amino-1-propanol in step 1 (e) and using phenol
30 in place of 4-fluorophenol in step 1 (h) afforded the title compound as yellow solid.

ES (+) MS m/e = 467 (MH⁺)

Example 9

35

(S)-1-(1-Hydroxy-3-phenylprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy-
pyrimidin-4-yl)imidazole

Following the procedures of examples 1 (e) - (h) except using (S)-2 amino-3-phenyl-1-propanol in place of (S)-2 amino-1-propanol in step 1 (e) and using phenol in place of 4-fluorophenol in step 1 (h) afforded the title compound as a yellow solid. ES (+) MS m/e = 467 (MH⁺)

5

Example 10

(R)-1-(1-Hydroxybut-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole

Following the procedures of examples 1 (e) - (h) except using (R)-2 amino-1-butanol in place of (S)-2 amino-1-propanol in step 1 (e) and using phenol in place of 4-fluorophenol in step 1 (h) afforded the title compound as white solid. ES (+) MS m/e = 405 (MH⁺)

10

Example 11

(S)-1-(1-Hydroxybut-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole

Following the procedures of examples 1 (e) - (h) except using (S)-2 amino-1-butanol in place of (S)-2 amino-1-propanol in step 1 (e) and using phenol in place of 4-fluorophenol in step 1 (h) afforded the title compound as white solid. ES (+) MS m/e = 405 (MH⁺)

20

Example 12

1-(1,3-Dihydroxyprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole

Following the procedures of examples 1 (e) - (h) except using 2-amino-1,3-propanediol in place of (S)-2 amino-1-propanol in step 1 (e) and using phenol in place of 4-fluorophenol in step 1 (h) afforded the title compound as a white solid. ES (+) MS m/e = 407 (MH⁺)

25

Example 13

(R)-1-(1-Hydroxy-2-phenyleth-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole

Following the procedures of examples 1 (e) - (h) except using (R)-2-amino-2-phenylethanol in place of (S)-2 amino-1-propanol in step 1 (e) and using phenol in place of 4-fluorophenol in step 1 (h) afforded the title compound as a white solid. ES (+) MS m/e = 453 (MH⁺)

30

35

Example-14

(S)-1-(1-Hydroxy-2-phenyleth-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy-pyrimidin-4-yl)imidazole

- 5 Following the procedures of examples 1 (e) - (h) except using (S)-2-amino-2-phenylethanol in place of (S)-2 amino-1-propanol in step 1 (e) and using phenol in place of 4-fluorophenol in step 1 (h) afforded the title compound as a white solid. ES (+) MS m/e = 453 (MH⁺)

Example-15

(S)-1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-chlorophenoxy)pyrimidin-4-yl]imidazole

- 10 Following the procedure of example 1 (h) except using 4-chlorophenol in place of 4-fluorophenol, afforded the title compound as a white solid.
15 ES (+) MS m/e = 425 (MH⁺)

Example 16

(R)-1-(1-Hydroxybut-2-yl)-4-(4-fluorophenyl)-5-[2-(4-chlorophenoxy)pyrimidin-4-yl]imidazole

- 20 Following the procedures of examples 1 (e) - (h) except using (R)-2 amino-1-butanol in place of (S)-2 amino-1-propanol in step 1 (e) and using 4-chlorophenol in place of 4-fluorophenol in step 1 (h) afforded the title compound as a white solid. ES (+) ES (+) MS m/e = 439 (MH⁺)

Example 17

(S)-1-(1-Hydroxybut-2-yl)-4-(4-fluorophenyl)-5-[2-(4-chlorophenoxy)pyrimidin-4-yl]imidazole

- 25 Following the procedures of examples 1 (e) - (h) except using (S)-2 amino-1-butanol in place of (S)-2 amino-1-propanol in step 1 (e) and using 4-chlorophenol in
30 place of 4-fluorophenol in step 1 (h) afforded the title compound as white solid. ES (+) MS m/e = 439 (MH⁺)

Example 18

(S)-1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-methylphenoxy)pyrimidin-4-yl]imidazole

- 35

Following the procedure of example 1 (h) except using 4-methylphenol in place of 4-fluorophenol, afforded the title compound as a white solid.

ES (+) MS m/e = 405 (MH⁺)

5

Example 19

(R)-1-(1-Hydroxybut-2-yl)-4-(4-fluorophenyl)-5-[2-(4-methylphenoxy)pyrimidin-4-yl]imidazole

Following the procedures of examples 1 (e) - (h) except using (R)-2 amino-1-butanol in place of (S)-2 amino-1-propanol in step 1 (e) and using 4-methylphenol in place of 4-fluorophenol in step 1 (h) afforded the title compound as white solid. ES (+) MS m/e = 419 (MH⁺)

10

Example 20

(S)-1-(1-Hydroxybut-2-yl)-4-(4-fluorophenyl)-5-[2-(4-methylphenoxy)pyrimidin-4-yl]imidazole

15

Following the procedures of examples 1 (e) - (h) except using (S)-2 amino-1-butanol in place of (S)-2 amino-1-propanol in step 1 (e) and using 4-methylphenol in place of 4-fluorophenol in step 1 (h) afforded the title compound as white solid. ES (+) MS m/e = 419 (MH⁺)

20

Example 21

1-(1,3-Dihydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-methylphenoxy)pyrimidin-4-yl]imidazole

25

Following the procedures of examples 1 (e) - (h) except using 2-amino-1,3-propanediol in place of (S)-2 amino-1-propanol in step 1 (e) and using 4-methylphenol in place of 4-fluorophenol in step 1 (h) afforded the title compound as a white solid. ES (+) MS m/e = 421 (MH⁺)

Example 22

(S)-1-(4-N-Morpholinyl-2-but-yl)-4-(4-fluorophenyl)-5-(2-phenoxy)pyrimidin-4-yl)imidazole

30

a) (S)-3-(t-butoxycarbonylamino)-1-butanol

35

Methyl (S)-3-(t-butoxycarbonylamino)-1-butyrate prepared according to (McIntosh and Acquah, Can. J. Chem **66**, 1752 (1988)) (3 g, 13.8 mmol) was dissolved in dry THF (100 mL) and cooled in an ice bath while stirring under argon. Lithium borohydride (305 mg, 13.9 mmol) was added and the mixture allowed to

warm to room temperature and stir for 16 hours. The solvent was evaporated and the residue taken up in EtOAc, washed with brine, dried (Na_2SO_4), and concentrated. The crude product was purified by flash chromatography (silica gel, 0-20% EtOAc/hexane) to give the title compound as a white solid. ES (+) MS $m/e = 190$ (MH^+)

5 b) (S)-3-(t-butoxycarbonylamino)-1-butanol mesylate

The product of example 22 (a) (1.06 g, 5.6 mmol) was dissolved in CH_2Cl_2 (10 ml) and cooled in an ice bath while stirring under argon. Triethylamine (0.667 g, 6.6 mmol) was added followed by the dropwise addition of methanesulfonyl chloride (0.687 g, 6 mmol). After 10 minutes the ice bath was removed and the
10 mixture stirred at rt for 2 h. The solvent was stripped off and the title compound was isolated as a yellow oil. ES (+) MS $m/e = 268$ (MH^+), 290 (MNa^+)

c) 4-[(S)-3-(t-butoxycarbonylamino)but-1-yl]morpholine

The crude product of exaple 22 (b) (~5.6 mmol) was dissolved in morpholine (10 mL) and heated to 57 °C while stirring under argon for 1 h. The solvent was
15 pumped off, and the residue was partitioned between EtOAc and water. The organic phase was washed with brine, dried (Na_2SO_4), and concentrated. The crude product was purified by flash chromatography (silica gel, 2-5% MeOH/ CH_2Cl_2) to give the title compound as a white solid. ES (+) MS $m/e = 259$ (MH^+)


d) 4-(3-aminobut-1-yl)morpholine

20 The product of example 22 (c) (0.41 g, 1.59 mmol) was dissolved in CH_2Cl_2 (10 mL) and cooled in an ice bath while stirring under argon. A solution of trifluoroacetic acid (5 mL) in CH_2Cl_2 (5 mL) was added and the mixture allowed to warm to room temperature for 1 h. The solvents were evaporated to give the title compound as an oil. ES (+) MS $m/e = 159$ (MH^+)


25 e) (S)-1-(4-N-Morpholinyl-2-but-yl)-4-(4-fluorophenyl)-5-(2-phenoxy-
pyrimidin-4-yl)imidazole

Following the procedures of examples 1 (e) - (h) except using the crude product of 22 (d) plus excess triethylamine in place of (S)-2 amino-1-propanol in step 1 (e) and using phenol in place of 4-fluorophenol in step 1 (h) afforded the title
30 compound as a pink solid ES (+) MS $m/e = 474$ (MH^+)


By analagous methods to those indicated above the following compounds may be prepared:




#	R1	R2
23	methyl	hydroxymethyl
24	ethyl	hydroxymethyl
25	isopropyl	hydroxymethyl
26	2-(methyl)propyl	hydroxymethyl
27	phenyl	hydroxymethyl
28	benzyl	hydroxymethyl
29	phenethyl	hydroxymethyl
30	methyl	dimethylaminomethyl
31	ethyl	dimethylaminomethyl
32	isopropyl	dimethylaminomethyl
33	2-(methyl)propyl	dimethylaminomethyl
34	phenyl	dimethylaminomethyl
35	benzyl	dimethylaminomethyl
36	phenethyl	dimethylaminomethyl
37	methyl	carboxy
38	ethyl	carboxy
39	isopropyl	carboxy
40	2-(methyl)propyl	carboxy
41	phenyl	carboxy
42	benzyl	carboxy
43	phenethyl	carboxy



#	R1	R2
44	methyl	-CONH ₂
45	methyl	-CONHMe
46	methyl	CONMe ₂
47	methyl	methoxymethyl
48	methyl	phenoxymethyl
49	methyl	benzyloxymethyl
50	methyl	carbamoylmethyl
51	methyl	carbamoylmethyl
52	methyl	aminomethyl
53	methyl	<i>N</i> -(methyl)aminomethyl
54	methyl	<i>N</i> -(dimethyl)aminomethyl
55	methyl	<i>N</i> -(benzyl)aminomethyl
56	methyl	acetamidomethyl
57	methyl	benzamidomethyl
58	methyl	methanesulfonamidomethyl
59	methyl	benzenesulfonamidomethyl
60	methyl	phenylmethylsulfonamidomethyl
61	methyl	<i>N'</i> -ethylureidomethyl
62	methyl	<i>N'</i> -phenylureidomethyl
63	methyl	-CH ₂ OCH ₂ CH ₂ Ph
64	methyl	-CONHPh
65	methyl	-CONHBn
66	methyl	-CH ₂ OCH ₂ (4-MePh)
67	methyl	-CH ₂ OCH ₂ (4-FPh)
68	methyl	-CH ₂ OCH ₂ (4-ClPh)
69	methyl	-CH ₂ OCH ₂ (3-ClPh)
70	methyl	-CH ₂ O(4-MePh)
71	methyl	-CH ₂ O(4-FPh)



#	R1	R2
72	methyl	-CH ₂ O(4-ClPh)
73	methyl	-CH ₂ O(3-ClPh)
74	ethyl	-CONH ₂
75	ethyl	-CONHMe
76	ethyl	CONMe ₂
77	ethyl	methoxymethyl
78	ethyl	phenoxymethyl
79	ethyl	benzyloxymethyl
80	ethyl	carbamoyl
81	ethyl	carbamoylmethyl
82	ethyl	aminomethyl
83	ethyl	<i>N</i> -(methyl)aminomethyl
84	ethyl	<i>N</i> -(dimethyl)aminomethyl
85	ethyl	<i>N</i> -(benzyl)aminomethyl
86	ethyl	acetamidomethyl
87	ethyl	benzamidomethyl
88	ethyl	methanesulfonamidomethyl
89	ethyl	benzenesulfonamidomethyl
90	ethyl	phenylmethylsulfonamidomethyl
91	ethyl	<i>N'</i> -ethylureidomethyl
92	ethyl	<i>N'</i> -phenylureidomethyl
93	ethyl	-CH ₂ OCH ₂ CH ₂ Ph
94	ethyl	-CONHPh
95	ethyl	-CONHBn
96	ethyl	-CH ₂ OCH ₂ (4-MePh)



97	ethyl	-CH ₂ OCH ₂ (4-FPh)
98	ethyl	-CH ₂ OCH ₂ (4-ClPh)
99	ethyl	-CH ₂ OCH ₂ (3-ClPh)
100	ethyl	-CH ₂ O(4-MePh)
101	ethyl	-CH ₂ O(4-FPh)
102	ethyl	-CH ₂ O(4-ClPh)
103	ethyl	-CH ₂ O(3-ClPh)


5 All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

10 The above description fully discloses the invention including preferred embodiments thereof. Modifications and improvements of the embodiments specifically disclosed herein are within the scope of the following claims. Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. Therefore, the

15 Examples herein are to be construed as merely illustrative and not a limitation of the scope of the present invention in any way. The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows.

What is Claimed is:

1. A compound represented by the formula :



5 wherein

- R₁ is 4-pyridyl, pyrimidinyl, 4-pyridazinyl, 1,2,4-triazin-5-yl, quinolyl, isoquinolinyl, or quinazolin-4-yl ring, which ring is substituted with Y-R_a and optionally with an additional independent substituent selected from C₁₋₄ alkyl, halogen, hydroxyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, C₁₋₄ alkylsulfinyl, CH₂OR₁₂, amino, mono and di- C₁₋₆ alkyl substituted amino, an N-heterocyclyl ring which
- 10 ring has from 5 to 7 members and optionally contains an additional heteroatom selected from oxygen, sulfur or NR₁₅, N(R₁₀)C(O)R_b or NHR_a;

Y is oxygen or sulfur;

- R₄ is phenyl, naphth-1-yl or naphth-2-yl, or a heteroaryl, which is optionally
- 15 substituted by one or two substituents, each of which is independently selected, and which, for a 4-phenyl, 4-naphth-1-yl, 5-naphth-2-yl or 6-naphth-2-yl substituent, is halogen, cyano, nitro, C(Z)NR₇R₁₇, C(Z)OR₁₆, (CR₁₀R₂₀)_vCOR₁₂, SR₅, SOR₅, OR₁₂, halo-substituted-C₁₋₄ alkyl, C₁₋₄ alkyl, ZC(Z)R₁₂, NR₁₀C(Z)R₁₆, or (CR₁₀R₂₀)_vNR₁₀R₂₀ and which, for other
- 20 positions of substitution, is halogen, cyano, C(Z)NR₁₃R₁₄, C(Z)OR₃, (CR₁₀R₂₀)_m"COR₃, S(O)_mR₃, OR₃, halo-substituted-C₁₋₄ alkyl, C₁₋₄ alkyl, (CR₁₀R₂₀)_m"NR₁₀C(Z)R₃, NR₁₀S(O)_m"R₈, NR₁₀S(O)_m"NR₇R₁₇, ZC(Z)R₃ or (CR₁₀R₂₀)_m"NR₁₃R₁₄;

Z is oxygen or sulfur;

- 25 n is an integer having a value of 1 to 10;
- m is 0, or the integer 1 or 2;
- m' is an integer having a value of 1 or 2,
- m" is 0, or an integer having a value of 1 to 5;
- v is 0, or an integer having a value of 1 or 2;
- 30 R₂ is -C(H) (A) (R₂₂);

A is an optionally substituted aryl, heterocyclyl, or heteroaryl ring, or A is a substituted C₁₋₁₀ alkyl;

- R₂₂ is an optionally substituted C₁₋₁₀ alkyl;
- R_a is aryl, arylC₁₋₆alkyl, heterocyclic, heterocyclylC₁₋₆ alkyl, heteroaryl, heteroarylC₁₋₆alkyl, wherein each of these moieties may be optionally substituted;
- 5 R_b is hydrogen, C₁₋₆ alkyl, C₃₋₇ cycloalkyl, aryl, arylC₁₋₄ alkyl, heteroaryl, heteroarylC₁₋₄alkyl, heterocyclyl, or heterocyclylC₁₋₄ alkyl, wherein each of these moieties may be optionally substituted;
- R₃ is heterocyclyl, heterocyclylC₁₋₁₀ alkyl or R₈;
- R₅ is hydrogen, C₁₋₄ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl or NR₇R₁₇, excluding the
10 moieties SR₅ being SNR₇R₁₇ and SOR₅ being SOH;
- R₆ is hydrogen, a pharmaceutically acceptable cation, C₁₋₁₀ alkyl, C₃₋₇ cycloalkyl, aryl, arylC₁₋₄ alkyl, heteroaryl, heteroarylC₁₋₄alkyl, heterocyclyl, aroyl, or C₁₋₁₀ alkanoyl;
- R₇ and R₁₇ is each independently selected from hydrogen or C₁₋₄ alkyl or R₇ and
15 R₁₇ together with the nitrogen to which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR₁₅;
- R₈ is C₁₋₁₀ alkyl, halo-substituted C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₃₋₇ cycloalkyl, C₅₋₇ cycloalkenyl, aryl, arylC₁₋₁₀ alkyl, heteroaryl, heteroarylC₁₋₁₀
20 alkyl, (CR₁₀R₂₀)_nOR₁₁, (CR₁₀R₂₀)_nS(O)_mR₁₈, (CR₁₀R₂₀)_nNHS(O)₂R₁₈, (CR₁₀R₂₀)_nNR₁₃R₁₄; wherein the aryl, arylalkyl, heteroaryl, heteroaryl alkyl may be optionally substituted;
- R₉ is hydrogen, C(Z)R₁₁ or optionally substituted C₁₋₁₀ alkyl, S(O)₂R₁₈, optionally substituted aryl or optionally substituted aryl-C₁₋₄ alkyl;
- 25 R₁₀ and R₂₀ is each independently selected from hydrogen or C₁₋₄ alkyl;
- R₁₁ is hydrogen, C₁₋₁₀ alkyl, C₃₋₇ cycloalkyl, heterocyclyl, heterocyclyl C₁₋₁₀alkyl, aryl, arylC₁₋₁₀ alkyl, heteroaryl or heteroarylC₁₋₁₀ alkyl, wherein these moieties may be optionally substituted;
- R₁₂ is hydrogen or R₁₆;
- 30 R₁₃ and R₁₄ is each independently selected from hydrogen or optionally substituted C₁₋₄ alkyl, optionally substituted aryl or optionally substituted aryl-C₁₋₄ alkyl, or together with the nitrogen which they are attached form a heterocyclic ring of 5 to 7 members which ring optionally contains an additional heteroatom selected from oxygen, sulfur or NR₉ ;
- 35 R₁₅ is R₁₀ or C(Z)-C₁₋₄ alkyl;

R₁₆ is C₁₋₄ alkyl, halo-substituted-C₁₋₄ alkyl, or C₃₋₇ cycloalkyl;
 R₁₈ is C₁₋₁₀ alkyl, C₃₋₇ cycloalkyl, heterocyclyl, aryl, aryl₁₋₁₀alkyl, heterocyclyl,
 heterocyclyl-C₁₋₁₀alkyl, heteroaryl or heteroaryl₁₋₁₀alkyl;
 or a pharmaceutically acceptable salt thereof.

- 5 2. The compound according to Claim 1 wherein R₁ is a substituted 4-pyridyl or 4-pyrimindyl.
3. The compound according to Claim 2 wherein the R_a moiety is optionally
 10 substituted aryl or arylalkyl.
4. The compound according to Claim 1 or 3 wherein the R_a moiety is
 substituted one or more times with halogen; C₁₋₄ alkyl; halosubstituted; hydroxy;
 hydroxy substituted C₁₋₄ alkyl; (CR₁₀R₂₀)_q C₁₋₄ alkoxy; (CR₁₀R₂₀)_q
 15 S(O)_malkyl; (CR₁₀R₂₀)_qS(O)_m aryl; (CR₁₀R₂₀)_qC(O)OR₁₁;
 (CR₁₀R₂₀)_qC(O)R₁₁; (CR₁₀R₂₀)_qOC(O)R_c;
 -O-(CH₂)_s-O-; (CR₁₀R₂₀)_qNR₁₃R₁₄; (CR₁₀R₂₀)_qN(R₁₀)C(O)R_b;
 (CR₁₀R₂₀)_qC(O)NR₁₃R₁₄; (CR₁₀R₂₀)_qC(O)NR₁₀R_c;
 (CR₁₀R₂₀)_qS(O)₂NR₁₃R₁₄; (CR₁₀R₂₀)_qS(O)₂ NR₁₀R_c;
 20 (CR₁₀R₂₀)_qN(R₁₀)S(O)₂ R_c; cyano, nitro, N-heterocyclyl ring which ring has from
 5 to 7 members and optionally contains an additional heteroatom selected from
 oxygen, sulfur or NR₁₅; aryl; arylalkyl; aryloxy; or arylalkyloxy; and wherein the
 aryl, alkylalkyl, aryloxy and arylalkyloxy may be optionally substituted themselves
 one to two times by halogen; hydroxy; hydroxy substituted alkyl; C₁₋₁₀ alkoxy;
 25 S(O)_m alkyl; amino, NR₇R₁₇ group; C₁₋₄ alkyl, or halosubstituted C₁₋₄ alkyl;
 q is 0 or an integer having a value of 1 to 4;
 R_b is hydrogen, C₁₋₆ alkyl, C₃₋₇ cycloalkyl, aryl, arylC₁₋₄ alkyl, heteroaryl,
 heteroarylC₁₋₄alkyl, heterocyclyl, or a heterocyclylC₁₋₄ alkyl moiety; and
 R_c is C₁₋₆ alkyl, C₃₋₇ cycloalkyl, aryl, arylC₁₋₄ alkyl, heteroaryl,
 30 heteroarylC₁₋₄alkyl, heterocyclyl, or a heterocyclylC₁₋₄ alkyl moiety, wherein all of
 these moieties may be optionally substituted.
5. The compound according to Claim 4 wherein the substituent group on the R_a
 moiety is halogen, halosubstituted alkyl, hydroxy, cyano, alkyl, aryl, alkoxy,

aryloxy, arylalkyloxy, alkylthio, alkylsulfonyl, aminocarbonyl, acetamido, propionamido, methylenedioxy, tetrazole, methyltetrazolyl.


6. The compound according to Claim 2 wherein R₄ is an optionally substituted
5 phenyl.
7. The compound according to Claim 6 wherein the phenyl is substituted one or more times independently by halogen, SR₅, S(O)R₅, OR₁₂, halo-substituted-C₁₋₄ alkyl, or C₁₋₄ alkyl.
10
8. The compound according to any of Claims 1 to 7 wherein A is an optionally substituted aryl, heteroaryl or heterocyclic ring.
9. The compound according to any of Claims 1 to 7 wherein A is a substituted
15 C₁₋₁₀ alkyl.
10. The compound according to Claim 9 wherein A is a hydroxy substituted C₁₋₁₀ alkyl.
11. The compound according to Claim 1 or 9 wherein R₂₂ is a C₁₋₁₀ alkyl or a
20 hydroxy substituted C₁₋₁₀ alkyl.
12. The compound according to Claim 1 which is:
1-(4-N-Morpholinyl-2-but-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-
25 yl)imidazole;
1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-fluorophenoxy)pyrimidin-4-yl]imidazole
1-(1,3-Dihydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-fluorophenoxy)pyrimidin-4-yl]imidazole
30 1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole
1-(1-Phenoxyprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole
1-(1-Hydroxy-3-phenylprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole
1-(1-Hydroxybut-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole

- 1-(1,3-Dihydroxyprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole
- 1-(1-Hydroxy-2-phenyleth-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole
- 5 1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-chlorophenoxy)pyrimidin-4-yl]imidazole
- 1-(1-Hydroxybut-2-yl)-4-(4-fluorophenyl)-5-[2-(4-chlorophenoxy)pyrimidin-4-yl]imidazole
- 1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-methylphenoxy)pyrimidin-4-yl]imidazole
- 10 1-(1-Hydroxybut-2-yl)-4-(4-fluorophenyl)-5-[2-(4-methylphenoxy)pyrimidin-4-yl]imidazole
- 1-(1,3-Dihydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-methylphenoxy)pyrimidin-4-yl]imidazole;
- 15 or a pharmaceutically acceptable salt thereof.
13. A pharmaceutical composition comprising an effective amount of a compound according to any of Claims 1 to 12 and a pharmaceutically acceptable carrier or diluent.
- 20 14. A method of treating inflammation in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of Formula (I) according to any of Claims 1 to 12.
- 25 15. A method of treating a CSBP/RK/p38 kinase mediated disease in a mammal in need thereof, which method comprises administering to said mammal an effective amount of a compound of Formula (I) according to any of Claims 1 to 12.
16. The method according to claim 15 wherein the CSBP/RK/p38 kinase
- 30 mediated disease is psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, gout, traumatic arthritis, rubella arthritis and acute synovitis, rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic condition, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, Alzheimer's disease, stroke, neurotrauma, asthma, adult respiratory distress
- 35 syndrome, cerebral malaria, chronic pulmonary inflammatory disease, silicosis,


pulmonary sarcososis, bone resorption disease, osteoporosis, restenosis, stroke, cardiac and renal reperfusion injury, chronic renal failure, congestive heart failure, angiogenic diseases, thrombosis, glomerularnephritis, diabetes, graft vs. host reaction, allograft rejection, inflammatory bowel disease, Crohn's disease, ulcerative
 5 colitis, multiple sclerosis, muscle degeneration, eczema, contact dermatitis, psoriasis, sunburn, or conjunctivitis.

17. A process for preparing a compound of Formula (I) as defined in Claim 1 which comprises reacting a compound of the Formula (II) :

10



with a compound of the Formula (III):



15

wherein p is 0 or 2; and a base strong enough to deprotonate the isonitrile moiety of Formula (II); and R₁, R₂ and R₄ are as defined in Claim 1 or are precursors of the groups R₁, R₂ and R₄ and Ar is an optionally substituted phenyl group, and thereafter if necessary, converting a precursor of R₁, R₂ and R₄ to a group R₁, R₂
 20 and R₄.

18. The process according to Claim 17 wherein p=0.

19. The process according to Claim 17 wherein p=2.

25

20. The process according to Claim 19 wherein the base is an amine, a carbonate, a hydride, or an alkyl or aryl lithium reagent.

21. The process according to Claim 19 wherein the imine of Formula (III), is
 30 isolated prior to reaction with Formula (II).


22. The process according to Claim 19 wherein the imine of Formula (III), is formed in situ prior to reaction with Formula (II).

23. The process according to Claim 22 wherein the imine is formed in situ by reacting an aldehyde of the formula R_1CHO , wherein R_1 is as defined for Formula (I), with a primary amine of the formula R_2NH_2 , wherein R_2 is as defined for Formula (I).

24. The process according to Claim 23 wherein formation of the imine in situ utilizes dehydrating conditions.


25. The process according to Claim 23 or 24 wherein the solvent is N,N-dimethylformamide (DMF), a halogenated solvent, tetrahydrofuran (THF), dimethylsulfoxide (DMSO), an alcohol, benzene, toluene, MeCN, or DME.

26. The process according to Claim 23 wherein the aldehyde R_1CHO is a pyrimidine aldehyde of the formula:



wherein X is YRa, and X_1 is defined as the optional substituent group on the R_1 moiety in Formula (I) according to Claim 1, to yield a compound of Formula (I) or a pharmaceutically acceptable salt thereof.


27. The process according to Claim 24 wherein the aldehyde R_1CHO is a pyridine aldehyde of the formula:



wherein X is YRa, and X_1 is defined as the optional substituent group on the R_1 moiety in Formula (I) according to Claim 1, to yield a compound of Formula (I) or a pharmaceutically acceptable salt thereof.


28. The process according to Claim 27 wherein the compound is:
- 1-(4-N-Morpholinyl-2-but-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole;
- 1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-fluorophenoxy)pyrimidin-4-yl]imidazole
- 5 1-(1,3-Dihydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-fluorophenoxy)pyrimidin-4-yl]imidazole
- 1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole
- 1-(1-Phenoxyprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole
- 10 1-(1-Hydroxy-3-phenylprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole
- 1-(1-Hydroxybut-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole
- 1-(1,3-Dihydroxyprop-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole
- 15 1-(1-Hydroxy-2-phenyleth-2-yl)-4-(4-fluorophenyl)-5-(2-phenoxy pyrimidin-4-yl)imidazole
- 1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-chlorophenoxy)pyrimidin-4-yl]imidazole
- 1-(1-Hydroxybut-2-yl)-4-(4-fluorophenyl)-5-[2-(4-chlorophenoxy)pyrimidin-4-yl]imidazole
- 20 1-(1-Hydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-methylphenoxy)pyrimidin-4-yl]imidazole
- 1-(1-Hydroxybut-2-yl)-4-(4-fluorophenyl)-5-[2-(4-methylphenoxy)pyrimidin-4-yl]imidazole
- 25 1-(1,3-Dihydroxyprop-2-yl)-4-(4-fluorophenyl)-5-[2-(4-methylphenoxy)pyrimidin-4-yl]imidazole
- or a pharmaceutically acceptable salt thereof.

29. A compound, or pharmaceutically acceptable salt thereof, which is:



R1	R2
methyl	hydroxymethyl
ethyl	hydroxymethyl
isopropyl	hydroxymethyl
2-(methyl)propyl	hydroxymethyl
phenyl	hydroxymethyl
benzyl	hydroxymethyl
phenethyl	hydroxymethyl
methyl	dimethylaminomethyl
ethyl	dimethylaminomethyl
isopropyl	dimethylaminomethyl
2-(methyl)propyl	dimethylaminomethyl
phenyl	dimethylaminomethyl
benzyl	dimethylaminomethyl
phenethyl	dimethylaminomethyl
methyl	carboxy
ethyl	carboxy
isopropyl	carboxy
2-(methyl)propyl	carboxy
phenyl	carboxy
benzyl	carboxy
phenethyl	carboxy
methyl	-CONH ₂
methyl	-CONHMe
methyl	-CONMe ₂

30. A compound, or pharmaceutically acceptable salt thereof, which is:



R1	R2
methyl	methoxymethyl
methyl	phenoxymethyl
methyl	benzyloxymethyl
methyl	carbamoyl
methyl	carbamoylmethyl
methyl	aminomethyl
methyl	<i>N</i> -(methyl)aminomethyl
methyl	<i>N</i> -(dimethyl)aminomethyl
methyl	<i>N</i> -(benzyl)aminomethyl
methyl	acetamidomethyl
methyl	benzamidomethyl
methyl	methanesulfonamidomethyl
methyl	benzenesulfonamidomethyl
methyl	phenylmethylsulfonamidomethyl
methyl	<i>N'</i> -ethylureidomethyl
methyl	<i>N'</i> -phenylureidomethyl
methyl	-CH ₂ OCH ₂ CH ₂ Ph
methyl	-CONHPh
methyl	-CONHBn
methyl	-CH ₂ OCH ₂ (4-MePh)
methyl	-CH ₂ OCH ₂ (4-FPh)
methyl	-CH ₂ OCH ₂ (4-ClPh)
methyl	-CH ₂ OCH ₂ (3-ClPh)
methyl	-CH ₂ O(4-MePh)
methyl	-CH ₂ O(4-FPh)
methyl	-CH ₂ O(4-ClPh)

methyl	-CH ₂ O(3-ClPh)
ethyl	-CONH ₂
ethyl	-CONHMe
ethyl	CONMe ₂
ethyl	methoxymethyl
ethyl	phenoxymethyl
ethyl	benzyloxymethyl
ethyl	carbamoylmethyl
ethyl	carbamoylmethyl
ethyl	aminomethyl
ethyl	<i>N</i> -(methyl)aminomethyl
ethyl	<i>N</i> -(dimethyl)aminomethyl
ethyl	<i>N</i> -(benzyl)aminomethyl
ethyl	acetamidomethyl
ethyl	benzamidomethyl
ethyl	methanesulfonamidomethyl
ethyl	benzenesulfonamidomethyl
ethyl	phenylmethylsulfonamidomethyl
ethyl	<i>N'</i> -ethylureidomethyl
ethyl	<i>N'</i> -phenylureidomethyl
ethyl	-CH ₂ OCH ₂ CH ₂ Ph
ethyl	-CONHPh
ethyl	-CONHBn
ethyl	-CH ₂ OCH ₂ (4-MePh)
ethyl	-CH ₂ OCH ₂ (4-FPh)
ethyl	-CH ₂ OCH ₂ (4-ClPh)
ethyl	-CH ₂ OCH ₂ (3-ClPh)
ethyl	-CH ₂ O(4-MePh)
ethyl	-CH ₂ O(4-FPh)
ethyl	-CH ₂ O(4-ClPh)
ethyl	-CH ₂ O(3-ClPh)

31. A pharmaceutical composition comprising a compound according to Claim 29 or 30, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or diluent.
- 5 32. A method of treating a CSBP/RK/p38 kinase mediated disease in a mammal in need thereof, which method comprises administering to said mammal an effective amount of a compound according to Claim 29 or 30.
- 10 33. The method according to claim 32 wherein the CSBP/RK/p38 kinase mediated disease is psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, gout, traumatic arthritis, rubella arthritis and acute synovitis, rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic condition, sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, Alzheimer's disease, stroke, neurotrauma, asthma, adult respiratory distress
- 15 syndrome, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcososis, bone resorption disease, osteoporosis, restenosis, stroke, cardiac and renal reperfusion injury, chronic renal failure, congestive heart failure, angiogenic diseases, thrombosis, glomerularnephritis, diabetes, graft vs. host reaction, allograft rejection, inflammatory bowel disease, Crohn's disease, ulcerative
- 20 colitis, multiple sclerosis, muscle degeneration, eczema, contact dermititis, psoriasis, sunburn, or conjunctivitis.
34. The process according to Claim 17 wherein the compound of formula (I) is defined in Claim 29 or 30.

INTERNATIONAL SEARCH REPORT

Int. national application No.

PCT/US98/13809

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/235.8, 242, 252, 259, 274, 307, 314, 341; 544/122, 182, 224, 284, 315; 546/148, 167, 274.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4,900,352 A (WADA et al.) 13 February 1990, the entire document.	1-3, 5-7, 12, 17-34
A	US 4,725,600 A (TAKAYA et al.) 16 February 1988, the entire document.	1-3, 5-7, 17-34
A	US 4,427,437 A (SERBAN et al.) 24 January 1984, the entire document.	1-3, 5-7, 17-34

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

27 AUGUST 1998

Date of mailing of the international search report

28 SEP 1998

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

YOGENDRA N. GUPTA

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/13809

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6): A61K 31/44, 31/47, 31/50, 31/505, 31/495, 31/53, 31/535; C07D 213/02, 215/00, 217/00, 253/06, 237/06, 239/34, 239/72

A. CLASSIFICATION OF SUBJECT MATTER:

US CL : 514/235.8, 242, 252, 259, 274, 307, 314, 341; 544/122, 182, 224, 284, 315; 546/148, 167, 274.1

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/13809

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☒ Claims Nos.: 4, 8-11, 13-16
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.